

Subtly Broken Symmetries

Domènec Espriu

Departament d'Estructura i Constituents de la Matèria

and

Institut de Ciències del Cosmos



UNIVERSITAT DE BARCELONA



With thanks to Joan Soto



Institut de Ciències del Cosmos







The Nobel Prize in Physics 2008

"for the discovery of the mechanism of spontaneous broken symmetry in subatomic physics"

Spontaneous breaking of a symmetry

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

- At high temperatures the system is disordered and unmagnetized

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

- At high temperatures the system is disordered and unmagnetized

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

- At high temperatures the system is disordered and unmagnetized : The symmetry of the hamiltonian ($s_i \rightarrow -s_i$) is respected
- Below a certain temperature T_c the system is magnetized

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

- At high temperatures the system is disordered and unmagnetized : The symmetry of the hamiltonian ($s_i \rightarrow -s_i$) is respected
- Below a certain temperature T_c the system is magnetized : The symmetry is spontaneously broken

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

- At high temperatures the system is disordered and unmagnetized : The symmetry of the hamiltonian ($s_i \rightarrow -s_i$) is respected
- Below a certain temperature T_c the system is magnetized : The symmetry is spontaneously broken
- Symmetry group of the problem: $G = Z_2 = \{1, -1\}$

Spontaneous breaking of a symmetry

- The simplest statistical-mechanics system is the Ising model (E.Ising, 1925)

$$H = -\beta J \sum_{i,j} s_i s_j, \quad J > 0$$

- At high temperatures the system is disordered and unmagnetized : The symmetry of the hamiltonian ($s_i \rightarrow -s_i$) is respected
- Below a certain temperature T_c the system is magnetized : The symmetry is spontaneously broken
- Symmetry group of the problem: $G = Z_2 = \{1, -1\}$
- Symmetry group of the solution: $H = 1$

Ferromagnetism

(W. Heisenberg, 1928)

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

Ferromagnetism

(W. Heisenberg, 1928)

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

- Invariant under rotations: $G = SO(3)$

Ferromagnetism

(W. Heisenberg, 1928)

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

- Invariant under rotations: $G = SO(3)$
- The state of minimum energy is reached when all \mathbf{S}_i point in the same direction, it does not matter which one:
 $H = SO(2)$

Ferromagnetism

(W. Heisenberg, 1928)

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

- Invariant under rotations: $G = SO(3)$
- The state of minimum energy is reached when all \mathbf{S}_i point in the same direction, it does not matter which one:
 $H = SO(2)$
- Spontaneous symmetry breaking in the Ising or classical Heisenberg models is not a quantum effect

Ferromagnetism

(W. Heisenberg, 1928)


$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

- Invariant under rotations: $G = SO(3)$
- The state of minimum energy is reached when all \mathbf{S}_i point in the same direction, it does not matter which one:
 $H = SO(2)$

- Spontaneous symmetry breaking in the Ising or classical Heisenberg models is not a quantum effect
- There are excitations whose energy is arbitrarily close to the ground state solution: magnons or spin waves

Nambu-Goldstone modes

Nambu (1960), Goldstone (1961)



Ferromagnetic quantum system

The vectors \mathbf{S}_i are now $su(2)$ operators

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

One must choose a direction to quantize (e.g. \hat{z})

Ferromagnetic quantum system

The vectors \mathbf{S}_i are now $su(2)$ operators

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

One must choose a direction to quantize (e.g. \hat{z})

- H Invariant under rotations: $G = SO(3)$

Ferromagnetic quantum system

The vectors \mathbf{S}_i are now $su(2)$ operators

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

One must choose a direction to quantize (e.g. \hat{z})

- H Invariant under rotations: $G = SO(3)$
- If $J > 0$ the state of minimum energy is reached when all \mathbf{S}_i are either in the $|+\rangle$ or $|-\rangle$ state w.r.t. \hat{z} :
 $H = SO(2)$

Ferromagnetic quantum system

The vectors \mathbf{S}_i are now $su(2)$ operators

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

One must choose a direction to quantize (e.g. \hat{z})

- H Invariant under rotations: $G = SO(3)$
- If $J > 0$ the state of minimum energy is reached when all \mathbf{S}_i are either in the $|+\rangle$ or $|-\rangle$ state w.r.t. \hat{z} :

$$H = SO(2)$$

- Spontaneous symmetry breaking in the quantum Heisenberg models is now a quantum effect, but QM also imposes restrictions: no long range order for $d < 2$

Ferromagnetic quantum system

The vectors \mathbf{S}_i are now $su(2)$ operators

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad J > 0$$

One must choose a direction to quantize (e.g. \hat{z})

- H Invariant under rotations: $G = SO(3)$
- If $J > 0$ the state of minimum energy is reached when all \mathbf{S}_i are either in the $|+\rangle$ or $|-\rangle$ state w.r.t. \hat{z} :

$$H = SO(2)$$

- Spontaneous symmetry breaking in the quantum Heisenberg models is now a quantum effect, but QM also imposes restrictions: no long range order for $d < 2$
- There excitations whose energy is arbitrarily close to the ground state are now quantum states

Antiferromagnetism

If in

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j$$

we take the opposite sign for J , i.e. $J < 0$

Antiferromagnetism

If in

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

we take the opposite sign for J , i.e. $J < 0$

- The symmetry of the hamiltonian is still: $G = SO(3)$

Antiferromagnetism

If in

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

we take the opposite sign for J , i.e. $J < 0$

- The symmetry of the hamiltonian is still: $G = SO(3)$
- The ground state is the Néel (1930) state:

$$| \uparrow \uparrow \downarrow \downarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \dots \rangle$$

with respect to some arbitrary axis.

Antiferromagnetism

If in

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

we take the opposite sign for J , i.e. $J < 0$

- The symmetry of the hamiltonian is still: $G = SO(3)$
- The ground state is the Néel (1930) state:

$$| \uparrow \uparrow \downarrow \downarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \dots \rangle$$

with respect to some arbitrary axis.

- $\mathbf{S} = 0$, however an order parameter still exists $\sum_i (-1)^i \mathbf{S}_i$

Antiferromagnetism

If in

$$H = -\beta J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

we take the opposite sign for J , i.e. $J < 0$

- The symmetry of the hamiltonian is still: $G = SO(3)$
- The ground state is the Néel (1930) state:

$$| \uparrow \uparrow \downarrow \downarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \dots \rangle$$

with respect to some arbitrary axis.

- $\mathbf{S} = 0$, however an order parameter still exists $\sum_i (-1)^i \mathbf{S}_i$
- Unbroken group: $H = SO(2)$

Nambu-Goldstone states

The number of Nambu-Goldstone modes depends crucially
on two things

Nambu-Goldstone states

The number of Nambu-Goldstone modes depends crucially on two things

- The number of broken generators in the process $G \rightarrow H$

Nambu-Goldstone states

The number of Nambu-Goldstone modes depends crucially on two things

- The number of broken generators in the process $G \rightarrow H$
- Relativistic invariance

Nambu-Goldstone states

The number of Nambu-Goldstone modes depends crucially on two things

- The number of broken generators in the process $G \rightarrow H$
- Relativistic invariance

The ferromagnetic Heisenberg model has two broken generators, but is described by an effective hamiltonian that leads to first-order (Schrödinger) equation.

- The two degrees of freedom combine in the description of a single quasi-particle.

Nambu-Goldstone states

The number of Nambu-Goldstone modes depends crucially on two things

- The number of broken generators in the process $G \rightarrow H$
- Relativistic invariance

The ferromagnetic Heisenberg model has two broken generators, but is described by an effective hamiltonian that leads to first-order (Schrödinger) equation.

- The two degrees of freedom combine in the description of a single quasi-particle.

The antiferromagnetic Heisenberg model, also with two broken generators, is described by an effective wave equation, which is real ($c \rightarrow v_s$).

- The two broken generators correspond to two different quasi-particles

Superconductivity

Bardeen, Cooper, Schrieffer, 1956



UNIVERSITAT DE BARCELONA
UB



Institut de Ciències del Cosmos

Superconductivity

Bardeen, Cooper, Schrieffer, 1956

- H is invariant under local gauge transformations:
 $G = U(1)$. In today's parlance H is called the 'abelian Higgs model'.

Superconductivity

Bardeen, Cooper, Schrieffer, 1956

- H is invariant under local gauge transformations:
 $G = U(1)$. In today's parlance H is called the 'abelian Higgs model'.
- The ground state ("the minimum") is a superposition of Copper pairs, without a well-defined charge: $H = 1$

Superconductivity

Bardeen, Cooper, Schrieffer, 1956

- H is invariant under local gauge transformations:
 $G = U(1)$. In today's parlance H is called the 'abelian Higgs model'.
- The ground state ("the minimum") is a superposition of Cooper pairs, without a well-defined charge: $H = 1$
- The spontaneous breaking of the gauge symmetry is a purely quantum effect

Superconductivity

Bardeen, Cooper, Schrieffer, 1956

- H is invariant under local gauge transformations: $G = U(1)$. In today's parlance H is called the 'abelian Higgs model'.
- The ground state ("the minimum") is a superposition of Cooper pairs, without a well-defined charge: $H = 1$
- The spontaneous breaking of the gauge symmetry is a purely quantum effect
- The symmetry group G still exists but it is realized in a non-linear manner, Nambu, 1960
- Because the symmetry is local, there are no Goldstone bosons but the photon becomes massive (Meissner effect), Englert, Brout; Higgs, 1964

Superconductivity

Bardeen, Cooper, Schrieffer, 1956

- H is invariant under local gauge transformations: $G = U(1)$. In today's parlance H is called the 'abelian Higgs model'.
- The ground state ("the minimum") is a superposition of Cooper pairs, without a well-defined charge: $H = 1$
- The spontaneous breaking of the gauge symmetry is a purely quantum effect
- The symmetry group G still exists but it is realized in a non-linear manner, Nambu, 1960
- Because the symmetry is local, there are no Goldstone bosons but the photon becomes massive (Meissner effect), Englert, Brout; Higgs, 1964

Superconductivity

Autobiography in *Broken Symmetry: Selected Papers of Y. Nambu*, World Scientific

One day before publication of the BCS paper, Bob Schrieffer, still a student, came to Chicago to give a seminar on the BCS theory in progress. . . . I was very much disturbed by the fact that their wave function did not conserve electron number. It did not make sense. . . . At the same time I was impressed by their boldness and tried to understand the problem.

6



UNIVERSITAT DE BARCELONA



Institut de Ciències del Cosmos

I will begin by a short story about my background. I studied physics at the University of Tokyo. I was attracted to particle physics because of the three famous names, Nishina, Tomonaga and Yukawa, who were the founders of particle physics in Japan. But these people were at different institutions than mine. On the other hand, condensed matter physics was pretty good at Tokyo. I got into particle physics only when I came back to Tokyo after the war. In hindsight, though, I must say that my early exposure to condensed matter physics has been quite beneficial to me.

History repeats itself

1960 Midwest Conference in Theoretical Physics, Purdue University

A 'SUPERCONDUCTOR' MODEL OF ELEMENTARY PARTICLES AND ITS CONSEQUENCES by Y. Nambu (University of Chicago)[†]

(In absence of the author the paper was presented by G. Jona-Lasinio.)

1

In recent years it has become fashionable to apply field-theoretical techniques to the many-body problems one encounters in solid state physics and nuclear physics. This is not surprising because in a quantized field theory there is always the possibility of pair creation (real or virtual), which is essentially a many-body problem. We are familiar with a number of close analogies between ideas and problems in elementary particle theory and the corresponding ones in solid state physics. For example, the Fermi sea of electrons in a metal is analogous to the Dirac sea of electrons in the vacuum, and we speak about electrons and holes in both cases. Some people must have thought of the meson field as something like the shielded Coulomb field. Of course, in elementary particles we have more symmetries and invariance properties than in the other, and blind analogies are often dangerous.

What about elementary particles?

What about elementary particles?

- If we think of the vacuum state, the state with no particles, as a state in which there is nothing, it is hard to imagine..

What about elementary particles?

- If we think of the vacuum state, the state with no particles, as a state in which there is nothing, it is hard to imagine..
- But if the vacuum is just the ground state of a quantum mechanical system, why not?

Nambu, 1960

Linear sigma model

Gell-Mann, Levy, 1960



UNIVERSITAT DE BARCELONA
U B



Institut de Ciències del Cosmos

Linear sigma model

Gell-Mann, Levy, 1960

$$L = i\bar{\psi}_L \not{\partial} \psi_L + i\bar{\psi}_R \not{\partial} \psi_R - g\bar{\psi}_L \Sigma \psi_R - g\bar{\psi}_R \sigma^\dagger \psi_L + V(\Sigma^\dagger \Sigma)$$

$$\Sigma \simeq \sigma I + i\pi^a \tau^a$$

Linear sigma model

Gell-Mann, Levy, 1960

$$L = i\bar{\psi}_L \not{\partial} \psi_L + i\bar{\psi}_R \not{\partial} \psi_R - g\bar{\psi}_L \Sigma \psi_R - g\bar{\psi}_R \sigma^\dagger \psi_L + V(\Sigma^\dagger \Sigma)$$

$$\Sigma = \sigma I + i\pi^a \tau^a$$

- The model is invariant under $G = SU(2)_L \times SU(2)_R$

Linear sigma model

Gell-Mann, Levy, 1960

$$L = i\bar{\psi}_L \not{\partial} \psi_L + i\bar{\psi}_R \not{\partial} \psi_R - g\bar{\psi}_L \Sigma \psi_R - g\bar{\psi}_R \sigma^\dagger \psi_L + V(\Sigma^\dagger \Sigma)$$

$$\Sigma = \sigma I + i\pi^a \tau^a$$

• The model is invariant under $G = SU(2)_L \times SU(2)_R$

• $\psi_L \rightarrow L\psi_L, \quad \psi_R \rightarrow R\psi_R, \quad \Sigma \rightarrow L\Sigma R^\dagger$

Linear sigma model

Gell-Mann, Levy, 1960

$$L = i\bar{\psi}_L \not{\partial} \psi_L + i\bar{\psi}_R \not{\partial} \psi_R - g\bar{\psi}_L \Sigma \psi_R - g\bar{\psi}_R \sigma^\dagger \psi_L + V(\Sigma^\dagger \Sigma)$$

$$\Sigma = \sigma I + i\pi^a \tau^a$$

- The model is invariant under $G = SU(2)_L \times SU(2)_R$
- $\psi_L \rightarrow L\psi_L, \quad \psi_R \rightarrow R\psi_R, \quad \Sigma \rightarrow L\Sigma R^\dagger$
- By choosing V appropriately one finds $\Sigma = \Sigma_0 \neq 0$. Any value of Σ_0 is equivalent, e.g. $\Sigma_0 = I \Rightarrow H = SU(2)_V$

Linear sigma model

Gell-Mann, Levy, 1960

$$L = i\bar{\psi}_L \not{\partial} \psi_L + i\bar{\psi}_R \not{\partial} \psi_R - g\bar{\psi}_L \Sigma \psi_R - g\bar{\psi}_R \sigma^\dagger \psi_L + V(\Sigma^\dagger \Sigma)$$

$$\Sigma = \sigma I + i\pi^a \tau^a$$

- The model is invariant under $G = SU(2)_L \times SU(2)_R$
- $\psi_L \rightarrow L\psi_L, \quad \psi_R \rightarrow R\psi_R, \quad \Sigma \rightarrow L\Sigma R^\dagger$
- By choosing V appropriately one finds $\Sigma = \Sigma_0 \neq 0$. Any value of Σ_0 is equivalent, e.g. $\Sigma_0 = I \Rightarrow H = SU(2)_V$
- The original symmetry group G is realized in a non-linear manner (Nambu, 1960)
- $\Sigma = \sigma U, \quad U = e^{i\tilde{\pi}/f_\pi}$ and G acts as $U \rightarrow LUR^\dagger$

Linear sigma model

Gell-Mann, Levy, 1960

$$L = i\bar{\psi}_L \not{\partial} \psi_L + i\bar{\psi}_R \not{\partial} \psi_R - g\bar{\psi}_L \Sigma \psi_R - g\bar{\psi}_R \sigma^\dagger \psi_L + V(\Sigma^\dagger \Sigma)$$

$$\Sigma = \sigma I + i\pi^a \tau^a$$

- The model is invariant under $G = SU(2)_L \times SU(2)_R$
- $\psi_L \rightarrow L\psi_L, \quad \psi_R \rightarrow R\psi_R, \quad \Sigma \rightarrow L\Sigma R^\dagger$
- By choosing V appropriately one finds $\Sigma = \Sigma_0 \neq 0$. Any value of Σ_0 is equivalent, e.g. $\Sigma_0 = I \Rightarrow H = SU(2)_V$
- The original symmetry group G is realized in a non-linear manner (Nambu, 1960)
- $\Sigma = \sigma U, \quad U = e^{i\tilde{\pi}/f_\pi}$ and G acts as $U \rightarrow LUR^\dagger$

The Nambu-Jona-Lasinio model I

Nambu, Jona-Lasinio, 1961



UNIVERSITAT DE BARCELONA



Institut de Ciències del Cosmos

The Nambu-Jona-Lasinio model I

Nambu, Jona-Lasinio, 1961

- A model of relativistic fermions without mass with contact interactions

The Nambu-Jona-Lasinio model I

Nambu, Jona-Lasinio, 1961

- A model of relativistic fermions without mass with contact interactions
- The model is invariant under a global chiral symmetry $U_L(1) \otimes U_R(1)$

The Nambu-Jona-Lasinio model I

Nambu, Jona-Lasinio, 1961

- A model of relativistic fermions without mass with contact interactions
- The model is invariant under a global chiral symmetry $U_L(1) \otimes U_R(1)$
- The ground state, “the vacuum” is only invariant under the diagonal $U(1)$!

The Nambu-Jona-Lasinio model I

Nambu, Jona-Lasinio, 1961

- A model of relativistic fermions without mass with contact interactions
- The model is invariant under a global chiral symmetry $U_L(1) \otimes U_R(1)$
- The ground state, “the vacuum” is only invariant under the diagonal $U(1)$!
- Massless bosons appear: Nambu-Goldstone bosons

The Nambu-Jona-Lasinio model I

Nambu, Jona-Lasinio, 1961

- A model of relativistic fermions without mass with contact interactions
- The model is invariant under a global chiral symmetry $U_L(1) \otimes U_R(1)$
- The ground state, “the vacuum” is only invariant under the diagonal $U(1)$!
- Massless bosons appear: Nambu-Goldstone bosons
- Fermions become massive: dynamical mass generation

The Nambu–Jona-Lasinio (NJL) model

Y. Nambu, G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961)

The Lagrangian of the model is

$$L = -\bar{\psi}\gamma_{\mu}\partial_{\mu}\psi + g [(\bar{\psi}\psi)^2 - (\bar{\psi}\gamma_5\psi)^2]$$

It is invariant under ordinary and γ_5 gauge transformations

$$\begin{array}{ll} \psi \rightarrow e^{i\alpha}\psi, & \bar{\psi} \rightarrow \bar{\psi}e^{-i\alpha} \\ \psi \rightarrow e^{i\alpha\gamma_5}\psi, & \bar{\psi} \rightarrow \bar{\psi}e^{i\alpha\gamma_5} \end{array}$$

Mass equation

$$\frac{2\pi^2}{g\Lambda^2} = 1 - \frac{m^2}{\Lambda^2} \ln \left(1 + \frac{\Lambda^2}{m^2} \right)$$

where Λ is the invariant cut-off

Spectrum of bound states

nucleon number	mass μ	spin-parity	spectroscopic notation
0	0	0^-	1S_0
0	$2m$	0^+	3P_0
0	$\mu^2 > \frac{8}{3}m^2$	1^-	3P_1
± 2	$\mu^2 > 2m^2$	0^+	1S_0

The Nambu-Jona-Lasinio model II

Nambu, Jona-Lasinio, 1961



UNIVERSITAT DE BARCELONA
U B



Institut de Ciències del Cosmos

The Nambu-Jona-Lasinio model II

Nambu, Jona-Lasinio, 1961

- Massless relativistic fermions \rightarrow Massless nucleons

The Nambu-Jona-Lasinio model II

Nambu, Jona-Lasinio, 1961

- Massless relativistic fermions \rightarrow Massless nucleons
- $U_L(1) \otimes U(1)_R \rightarrow SU_L(2) \otimes SU_R(2)$

The Nambu-Jona-Lasinio model II

Nambu, Jona-Lasinio, 1961

- Massless relativistic fermions \rightarrow Massless nucleons
- $U_L(1) \otimes U(1)_R \rightarrow SU_L(2) \otimes SU_R(2)$
- The ground state is invariant only under diagonal $SU(2)$ isospin

The Nambu-Jona-Lasinio model II

Nambu, Jona-Lasinio, 1961

- Massless relativistic fermions \rightarrow Massless nucleons
- $U_L(1) \otimes U(1)_R \rightarrow SU_L(2) \otimes SU_R(2)$
- The ground state is invariant only under diagonal $SU(2)$ isospin
- Massless bosons \rightarrow Pions

The Nambu-Jona-Lasinio model II

Nambu, Jona-Lasinio, 1961

- Massless relativistic fermions \rightarrow Massless nucleons
- $U_L(1) \otimes U(1)_R \rightarrow SU_L(2) \otimes SU_R(2)$
- The ground state is invariant only under diagonal $SU(2)$ isospin
- Massless bosons \rightarrow Pions
- Nucleons become eventually massless

Nambu-Jona-Lasinio model?



Nambu-Jona-Lasinio model?

- A finite cut-off must be introduced

Nambu-Jona-Lasinio model?

- A finite cut-off must be introduced
- “*We are driven to the conclusion that the Hamiltonian method for strong interaction is dead and must be buried, although of course with deserved honor.*” (Landau, 1960)

Nambu-Jona-Lasinio model?

- A finite cut-off must be introduced
- “*We are driven to the conclusion that the Hamiltonian method for strong interaction is dead and must be buried, although of course with deserved honor.*” . (Landau, 1960)
- Nambu emphasized that the need of the cut-off only meant that high energy effects were unknown, but that the conclusions regarding the spectrum were firm

Nambu-Jona-Lasinio model?

- A finite cut-off must be introduced
- “We are driven to the conclusion that the Hamiltonian method for strong interaction is dead and must be buried, although of course with deserved honor.” (Landau, 1960)
- Nambu emphasized that the need of the cut-off only meant that high energy effects were unknown, but that the conclusions regarding the spectrum were firm
- In two dimensions and in the limit where the number of nucleons is $\rightarrow \infty$, the model is fully renormalizable and a bona fide QFT (Gross, Neveu, 1974)

QCD (~ 1972)



QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)

QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)
- The lagrangian is expressed in terms of quarks (Gell-Mann; Zweig, 1964) and gluons

QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)
- The lagrangian is expressed in terms of quarks (Gell-Mann; Zweig, 1964) and gluons
- However, only hadrons are physical

QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)
- The lagrangian is expressed in terms of quarks (Gell-Mann; Zweig, 1964) and gluons
- However, only hadrons are physical
- Analogies with NJL:

QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)
- The lagrangian is expressed in terms of quarks (Gell-Mann; Zweig, 1964) and gluons
- However, only hadrons are physical
- Analogies with NJL:
 - Dynamic mass generation

QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)
- The lagrangian is expressed in terms of quarks (Gell-Mann; Zweig, 1964) and gluons
- However, only hadrons are physical
- Analogies with NJL:
 - Dynamic mass generation
 - Spontaneous breakdown of chiral symmetry

QCD (~ 1972)

- Non-abelian Yang-Mills theory based on the group $SU(3)$ of color (Han, Nambu, 1965)
- The lagrangian is expressed in terms of quarks (Gell-Mann; Zweig, 1964) and gluons
- However, only hadrons are physical
- Analogies with NJL:
 - Dynamic mass generation
 - Spontaneous breakdown of chiral symmetry
 - Pions are (quasi-) Nambu-Goldstone bosons. Its mass is due to the soft breaking induced by current quark masses

QCD



Institut de Ciències del Cosmos

QCD

- Differences with NJL:

QCD


- Differences with NJL:
- Hadrons -in particular pions- are bound states of quarks, not nucleons

QCD

- Differences with NJL:
 - Hadrons -in particular pions- are bound states of quarks, not nucleons
 - NJL can nevertheless be reformulated in term of quarks, and is still widely used as a phenomenological low-energy approximation to QCD. It has however numerous shortcomings.

QCD

- Differences with NJL:
 - Hadrons -in particular pions- are bound states of quarks, not nucleons
 - NJL can nevertheless be reformulated in term of quarks, and is still widely used as a phenomenological low-energy approximation to QCD. It has however numerous shortcomings.
- Physical results do not depend on the cut-off thanks to the renormalizability of Yang-Mills theories (see below...)




Pions, (quasi-) Nambu-Goldstone bosons?

Pions, (quasi-) Nambu-Goldstone bosons?

This can nowadays be checked in lattice QCD by varying the bare quark mass:

Pions, (quasi-) Nambu-Goldstone bosons?

This can nowadays be checked in lattice QCD by varying the bare quark mass:



Giusti, 2006

Nambu's comment

Y. Nambu, preliminary Notes for the Nobel Lecture

In hindsight I regret that I should have explored in more detail the general mechanism of mass generation for the gauge field. But I thought the plasma and the Meissner effect had already established it. I also should have paid more attention to the Ginzburg-Landau theory which was a forerunner of the present Higgs description.

Someone we knew well was in Nambu's group ...

Someone we knew well was in Nambu's group ...



Someone we knew well was in Nambu's group ...



Pere Pascual de Sans
Sevilla, 1934 - Barcelona, 2006

Someone we knew well ...

Self-consistent models of strong interactions with chiral symmetry

Y. Nambu, P. Pascual, Nuovo Cimento 30 (1963) 354

Someone we knew well ...

Self-consistent models of strong interactions with chiral symmetry
Y. Nambu, P. Pascual, Nuovo Cimento 30 (1963) 354

“I was in Chicago for two years (1961-62) and Nambu gave to me two articles to read”



Someone we knew well ...

Self-consistent models of strong interactions with chiral symmetry
Y. Nambu, P. Pascual, Nuovo Cimento 30 (1963) 354

“I was in Chicago for two years (1961-62) and Nambu gave to me two articles to read”

- *Conservation of Isotopic Spin and Isotopic Gauge Invariance*,
Chen-Ning Yang, Robert L. Mills, Phys. Rev. 96, 191
(1954)

Someone we knew well ...

Self-consistent models of strong interactions with chiral symmetry
Y. Nambu, P. Pascual, Nuovo Cimento 30 (1963) 354

“I was in Chicago for two years (1961-62) and Nambu gave to me two articles to read”

- *Conservation of Isotopic Spin and Isotopic Gauge Invariance*,
Chen-Ning Yang, Robert L. Mills, Phys. Rev. 96, 191 (1954)
- *Quantum electrodynamics at small distances*, Murray
Gell-Mann, F.E. Low, Phys. Rev. 95, 1300 (1954)

Someone we knew well ...

Self-consistent models of strong interactions with chiral symmetry
Y. Nambu, P. Pascual, Nuovo Cimento 30 (1963) 354

“I was in Chicago for two years (1961-62) and Nambu gave to me two articles to read”

- *Conservation of Isotopic Spin and Isotopic Gauge Invariance*,
Chen-Ning Yang, Robert L. Mills, Phys. Rev. 96, 191
(1954)

- *Quantum electrodynamics at small distances*, Murray
Gell-Mann, F.E. Low, Phys. Rev. 95, 1300 (1954)

... both of them eventually proved to be crucial papers in the history of particle physics



Regge analysis of pion-pion (and pion-kaon) scattering for energy $s^{1/2} > 1.4$ GeV

J. R. Peláez

*Departamento de Física Teórica, II
(Métodos Matemáticos),*

*Facultad de Ciencias Físicas,
Universidad Complutense de Madrid,
E-28040, Madrid, Spain.*

and

F. J. Ynduráin

*Departamento de Física Teórica, C-XI
Universidad Autónoma de Madrid,
Canto Blanco,
E-28049, Madrid, Spain.*

Abstract

We perform a detailed Regge analysis of NN , πN , KN , $\pi\pi$ and πK scattering. From it, we find expressions that represent the $\pi\pi$ scattering amplitudes with an accuracy of a few percent, for exchange of isospin zero, and $\sim 15\%$ for exchange of isospin 1, and this for energies $s^{1/2} > 1.4$ GeV and for momentum transfers $|t|^{1/2} \lesssim 0.4$ GeV. These Regge formulas are perfectly compatible with the low energy ($s^{1/2} \sim 1.4$ GeV) scattering amplitudes deduced from $\pi\pi$ phase shift analyses as well as with higher energy ($s^{1/2} \gtrsim 1.4$ GeV) experimental $\pi\pi$ cross sections. They are also compatible with NN , KN and πN experimental cross sections using factorization, a property that we check with great precision. This contrasts with results from current phase shift analyses of the $\pi\pi$ scattering amplitude which bear little resemblance to reality in the region $1.4 < s^{1/2} < 2$ GeV, as they are not well defined and increasingly violate a number of physical requirements when the energy grows. πK scattering is also considered, and we present a Regge analysis for these processes valid for energies $s^{1/2} > 1.7$ GeV.

As a byproduct of our analysis we present also a fit of NN , πN and KN cross sections valid from c.m. kinetic energy $E_{\text{kin}} \simeq 1$ GeV to multi TeV energies.

arXiv:hep-ph/0312187v2 9 Feb 2004

Experimental status of the $\pi\pi$ isoscalar S wave at low energy: $f_0(600)$ pole and scattering length

R. García-Martín and J. R. Peláez
*Departamento de Física Teórica, II
(Métodos Matemáticos),
Facultad de Ciencias Físicas,
Universidad Complutense de Madrid,
E-28040, Madrid, Spain*

and F. J. Ynduráin
*Departamento de Física Teórica, C-XI
Universidad Autónoma de Madrid,
Canto Blanco,
E-28049, Madrid, Spain.*

Abstract

The experimental results obtained in the last few years on kaon decays ($K \rightarrow 2\pi$ and, above all, K_{e4} decays) allow a reliable, model independent determination of low energy $\pi\pi$ scattering in the S0 wave. Using them and, eventually, other sets of data, it is possible to give a precise parametrization of the S0 wave as well as to find the scattering length and effective range parameter. One can also perform an extrapolation to the pole of the “ σ resonance” [$f_0(600)$]. We obtain the results

$$a_0^{(0)} = 0.233 \pm 0.013 M_\pi^{-1}, \quad b_0^{(0)} = 0.285 \pm 0.012 M_\pi^{-3}$$

and, for the σ pole,

$$M_\sigma = 484 \pm 17 \text{ MeV}, \quad \Gamma_\sigma/2 = 255 \pm 10 \text{ MeV}.$$

arXiv:hep-ph/0701025v3 26 Jul 2007

Conclusions



Conclusions

Nature is indeed subtle



UNIVERSITAT DE BARCELONA



Institut de Ciències del Cosmos

Conclusions

Nature is indeed subtle

Thanks Paco!