




ATLAS



# LHC Results (HIGGS)



Mario Martínez



Taller de Altas Energías  
TAE 2014  
(Spanish School for HEP)

15-26 September  
Benasque



**Courses**

*QFT, SM and EWSB* Wolfgang Hollik (MPI – Munich)  
*QCD, jets and Monte Carlo* Germán Rodrigo (IFIC – Valencia)  
*Lattice* Elna Güntz (U. Granada)  
*Statistics* Carlos Mañá (CIEMAT – Madrid)  
*LHC results: Higgs* Mario Martínez (IFAE – Barcelona)  
*LHC results: EW, top, jets* Roberto Tenchini (CERN)  
*LHC results: Exotics* Greg Landsberg (Brown U.)  
*LHC Future* Ignacio Redondo (CIEMAT – Madrid)  
*Flavour Physics* Frederic Teubert (CERN)  
*Neutrinos* Gabriela Barenboim (IFIC – Valencia)  
*Heavy ions* Konrad Tywoniuk (U. Barcelona)  
*Effective QFT and BSM* José R. Espinosa (IFAE – Barcelona)  
*Supersymmetry* Dominik Stockinger (IKTP – Dresden)  
*Dark Matter* David Cordero (IFT – Madrid)  
*Cosmic Rays* Sergio Navas (U. Granada)  
*Gamma Rays* Manel Martínez\* (IFAE – Barcelona)  
*Cosmology* Ramon Miquel (IFAE – Barcelona)

\* to be confirmed

**Special Talks**

José María Hernández (CIEMAT – Madrid) Computing in LHC  
 Carlos Pérez de los Heros (U. Uppsala) IceCube  
 Carmen García (IFIC – Valencia) Future Accelerators  
 Detector Demonstration

**Organizers**

Juan Alcaraz (CIEMAT – Madrid)  
 José I. Illana (U. Granada)


**Registration**  
<http://benasque.org/2014tae>  
 (deadline: 15 July 2014)

## TAE SCHOOL FOR HIGH ENERGY PHYSICS BENASQUE, SEPTEMBER 2014



# Outline for Today

- Mass vs BEH Mechanism
- The BEH Mechanism
- Higgs boson production and decays
- Pre-LHC Higgs Searches at Colliders
- Building Blocks for a Discovery
- Higgs Discovery (Golden Channels)
- Higgs  $J^{PC}$
- *Other (Silver) Channels*



*"This could be the discovery of the century. Depending, of course, on how far down it goes."*

*Disclaimer: completely unbalanced set of results from CMS and ATLAS*



# Tomorrow...

$J^{PC}$

- *Other (Silver) Channels*
- Detailed study on Couplings
- Higgs width
- Invisibly decaying Higgs
- Higgs and Vacuum Stability
- Hierarchy Problem & SUSY
- Search for other Higgs
- What to expect in 2015 -- ?







I. Newton

# Mass

$$F = M a$$





A. Einstein

$$E = M c^2$$

Mass as a form of energy

Ley de Gravitación Universal de Newton

$$F_G = \frac{G m_T m_L}{r_{TL}^2}$$



Protón (uud)

Only 2% of the proton mass is due to quarks...most of it is QCD confinement energy


# Lesson 0

*Do not go around saying:*

*“The Higgs mechanism explains the origin of the mass in the Universe...”*

*It is imply incorrect...*










# EWK Symmetry Breaking

BHE mechanism makes the small range and weak interaction (massive Ws and Z)

Same time allows for fermion masses




big bang






# A world without mass ?

$$a_0 = \frac{1}{\alpha_{em} m_e} \text{ (natural units)}$$




$$G_F \propto \frac{1}{M_W^2}$$



Without giving masses to fermions  
it would be impossible to form stable atoms

Light W ? → Faster reactions and a Cold Universe



# How to include the mass ?

$$-m\psi\psi$$

*These terms are not allowed by gauge invariance*





$$\mathcal{L} = (D_\mu \phi)^\dagger D^\mu \phi - \mathcal{V}(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

$$D_\mu \phi = \partial_\mu \phi - ie A_\mu \phi$$

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

$$\mathcal{V}(\phi) = \alpha \phi^\dagger \phi + \beta (\phi^\dagger \phi)^2$$

Peter Higgs

$$\alpha < 0, \beta > 0$$

# 1964....

VOLUME 13, NUMBER 9

PHYSICAL REVIEW LETTERS

31 AUGUST 1964

---

**BROKEN SYMMETRY AND THE MASS OF GAUGE VECTOR MESONS\***

**F. Englert and R. Brout**

Faculté des Sciences, Université Libre de Bruxelles, Bruxelles, Belgium

(Received 26 June 1964)

VOLUME 13, NUMBER 16

PHYSICAL REVIEW LETTERS

19 OCTOBER 1964

---

**BROKEN SYMMETRIES AND THE MASSES OF GAUGE BOSONS**

**Peter W. Higgs**

Tait Institute of Mathematical Physics, University of Edinburgh, Edinburgh, Scotland

(Received 31 August 1964)

VOLUME 13, NUMBER 20

PHYSICAL REVIEW LETTERS

16 NOVEMBER 1964

---

**GLOBAL CONSERVATION LAWS AND MASSLESS PARTICLES\***


**G. S. Guralnik,<sup>†</sup> C. R. Hagen,<sup>‡</sup> and T. W. B. Kibble**

Department of Physics, Imperial College, London, England

(Received 12 October 1964)

# The BEH Mechanism

$$SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$$



Introduce one complex scalar doublet of  $SU(2)_L$  with  $Y = 1/2$ :

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \longleftrightarrow \mathcal{L} = (D^\mu \phi)^\dagger D_\mu \phi - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2$$

where  $D_\mu \phi = (\partial_\mu - igA_\mu^a \tau^a - ig'Y_\phi B_\mu)$ , ( $\tau^a = \sigma^a/2$ ,  $a=1, 2, 3$ ).

The SM symmetry is spontaneously broken when  $\langle \phi \rangle$  is chosen to be (e.g.):


$$\langle \phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} \quad \text{with} \quad v = \left( \frac{-\mu^2}{\lambda} \right)^{1/2} \quad (\mu^2 < 0, \lambda > 0)$$

The gauge boson mass terms arise from:

$$\begin{aligned} (D^\mu \phi)^\dagger D_\mu \phi &\longrightarrow \dots + \frac{1}{8} (0 \ v) (gA_\mu^a \sigma^a + g' B_\mu) (gA^{b\mu} \sigma^b + g' B^\mu) \begin{pmatrix} 0 \\ v \end{pmatrix} + \dots \\ &\longrightarrow \dots + \frac{1}{2} \frac{v^2}{4} [g^2 (A_\mu^1)^2 + g^2 (A_\mu^2)^2 + (-gA_\mu^3 + g' B_\mu)^2] + \dots \end{aligned}$$



# The BEH Mechanism



And correspond to the weak gauge bosons:

$$W_{\mu}^{\pm} = \frac{1}{\sqrt{2}}(A_{\mu}^1 \pm A_{\mu}^2) \longrightarrow \boxed{M_W = g \frac{v}{2}}$$

$$Z_{\mu}^0 = \frac{1}{\sqrt{g^2 + g'^2}}(gA_{\mu}^3 - g'B_{\mu}) \longrightarrow \boxed{M_Z = \sqrt{g^2 + g'^2} \frac{v}{2}}$$


while the linear combination orthogonal to  $Z_{\mu}^0$  remains massless and corresponds to the photon field:

$$A_{\mu} = \frac{1}{\sqrt{g^2 + g'^2}}(g'A_{\mu}^3 + gB_{\mu}) \longrightarrow \boxed{M_A = 0}$$

$$\cos \theta_w = \frac{g}{\sqrt{g^2 + g'^2}}, \quad \sin \theta_w = \frac{g'}{\sqrt{g^2 + g'^2}}$$

the  $W$  and  $Z$  masses are related by:  $\boxed{M_W = M_Z \cos \theta_w}$

# The BEH Mechanism



The scalar sector becomes more transparent in the unitary gauge:

$$\phi(x) = \frac{e^{\frac{i}{v}\vec{\chi}(x)\cdot\vec{\tau}}}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix} \xrightarrow{SU(2)} \phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix}$$

after which the Lagrangian becomes

$$\mathcal{L} = \mu^2 H^2 - \lambda v H^3 - \frac{1}{4} H^4 = -\frac{1}{2} M_H^2 H^2 - \sqrt{\frac{\lambda}{2}} M_H H^3 - \frac{1}{4} \lambda H^4$$

Three degrees of freedom, the  $\chi^a(x)$  Goldstone bosons, have been reabsorbed into the longitudinal components of the  $W_\mu^\pm$  and  $Z_\mu^0$  weak gauge bosons. One real scalar field remains:


the Higgs boson, H, with mass  $M_H^2 = -2\mu^2 = 2\lambda v^2$

and self-couplings:

$$\begin{array}{c} \text{H} \\ \text{H} \end{array} \text{---} \text{H} = -3i \frac{M_H^2}{v}$$

$$\begin{array}{c} \text{H} \\ \text{H} \end{array} \text{---} \text{H} = -3i \frac{M_H^2}{v^2}$$

# The BEH Mechanism



From  $(D^\mu \phi)^\dagger D_\mu \phi \rightarrow$  Higgs-Gauge boson couplings:

$$\begin{array}{c}
 V^\mu \\
 \text{wavy line} \\
 \text{---} \\
 V^\nu \\
 \text{wavy line}
 \end{array}
 \text{---} H = 2i \frac{M_V^2}{v} g^{\mu\nu}$$

$$\begin{array}{c}
 V^\mu \\
 \text{wavy line} \\
 \text{---} \\
 V^\nu \\
 \text{wavy line}
 \end{array}
 \begin{array}{l}
 \text{---} H \\
 \text{---} H
 \end{array} = 2i \frac{M_V^2}{v^2} g^{\mu\nu}$$

**Notice:** The entire Higgs sector depends on only two parameters, e.g.

$M_H$  and  $v$

$v$  measured in  $\mu$ -decay:



$$v = (\sqrt{2}G_F)^{-1/2} = 246 \text{ GeV}$$

$\rightarrow$

SM Higgs Physics depends on  $M_H$





# Consistency of Gauge Interactions





This illustrates the consistency of the Gauge Interactions

# $W_L^+ W_L^- \rightarrow W_L^+ W_L^-$ scattering



To restore unitarity one needs to postulate a Higgs boson with mass below 800 GeV

Otherwise new physics should come at the TeV scale to rescue us



# Fermion Masses



$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \quad Y_\phi = +1$$

Most of the free parameters in the SM model come from here


$$\mathcal{L}_F = -\lambda_e \bar{L} \Phi e_R - \lambda_d \bar{Q} \Phi d_R - \lambda_u \bar{Q} \tilde{\Phi} u_R + h.c.$$

$$\begin{aligned} \mathcal{L}_F &= -\frac{1}{\sqrt{2}} \lambda_e (\bar{\nu}_e, \bar{e}_L) \begin{pmatrix} 0 \\ v + H \end{pmatrix} e_R + \dots \\ &= -\frac{1}{\sqrt{2}} \lambda_e (v + H) \bar{e}_L e_R + \dots \end{aligned}$$

$$m_e = \frac{\lambda_e v}{\sqrt{2}}, \quad m_u = \frac{\lambda_u v}{\sqrt{2}}, \quad m_d = \frac{\lambda_d v}{\sqrt{2}}$$




# Why such a mass hierarchy ?











# Higgs Couplings to SM



Couplings proportional to masses of particles  
 → This determines the phenomenology




and via loops..



# Higgs Production (LHC)




Decreasing cross section  $\longrightarrow$




# Higgs Production (LHC)

For a Higgs of 125 GeV



Decreasing cross section



# Higgs Program in a Glance

Channel categories	ggF 	VBF 	VH 	ttH 
$\gamma\gamma$	✓	✓	✓	✓
ZZ (IIII)	✓	✓	✓	✓
WW (IInI)	✓	✓	✓	✓
$\tau\tau$	✓	✓	✓	✓
bb	?????	✓	✓	✓
$Z\gamma$	✓	✓		
$\mu\mu$	✓	✓		
Invisible	✓	✓	✓	



# Beat the background


The discovery channels are subject of signal rate, mass resolution, and the capacity to beat the background

For example *the a priori* good channel for Higgs production


$gg \rightarrow H, H \rightarrow b(\text{anti})b$

gets killed by the huge underneath (non-resonant) QCD-driven process


$gg \rightarrow b(\text{anti})b$







# Want a Nobel?...find a resonance






# The Golden Channels



Irreducible background from di-photons



...and background from  $\gamma$ +jets and jet-jet with jets faking photon signals...





**A resonance out of a huge non-resonant background**  
**Key: good photon energy/momentum reconstruction**




# The Golden Channels

Background dominated by  
Diboson ( $ZZ^*$ ) w/wo resolved Zs

with contributions from  
other processes (Z+jets) and top  
production with jets faking leptons



Very clean and high-resolution channel  
(few events piling up in a given mass will be  
enough to claim discovery)





# Pre-LHC Higgs Results



*Electroweak Fits*

*Tevatron Searches*

*Searches at LEP*


# Absolute limits on Higgs from LEP

LEP : e+ e- collider @ CERN : up to 210 GeV



Hot days at CERN ... by now we know it was nothing..

Excludes @ 95 % CL  
 $M_h < 114$  GeV





**KEEP  
CALM  
ITS A  
FALSE  
ALARM**

**A hard decision was taken to stop LEP and allow for the LHC program to start .. Quite the right decision.. we know now**



# Tevatron

Chicago



$$\sqrt{s} = 1.96 \text{ TeV}$$



Booster


CDF

DØ


Tevatron

p-bar source

Main Injector  
(new)










# Electroweak Fits

(indirect Higgs Mass constrains)



$$m_W^2 \left( 1 - \frac{m_W^2}{m_Z^2} \right) = \frac{\pi\alpha}{\sqrt{2}G_F} (1 + \Delta r)$$




$$\Delta r \sim \ln(m_H)$$


$$\Delta r \sim m_{top}^2$$

..as presented in  
Lepton Photon 2007

$$M_H < 160 \text{ GeV} / c^2$$




$qq \rightarrow WH, ZH$



# Tevatron in July 2010



Tevatron Run II Preliminary,  $\langle L \rangle = 5.9 \text{ fb}^{-1}$



If goes below 1 you exclude the SM Higgs for a given mass



# CERN (Geneva) LHC across the France-Switzerland border

## Approved in 1994



27 Km

1232 high-tech superconducting dipole magnets  
(at 1.8 K...the coldest **(and coolest)** place in the universe)

proton – proton 7-8 TeV in Run I (2010 – 2012)  
(13 TeV in Run II) (2015 -- )



# LHC construction









# LHC Construction





# Overall view of the LHC experiments.

LHC will run for 15 years..  
A total of  $6 \times 10^{15}$  collisions




CMS  
Point 5


LHC - B  
Point 8

CERN  
ATLAS  
Point 1

ALICE  
Point 2








**CMS**








ATLAS






**ATLAS**  
**design and construction**  
**over 15 years**  
**(this will take 1 minute)**











# ATLAS

(relevant to photon ID)





**Inner Detector** - Barrel (B) & End-cap (E) in 2T solenoidal magnetic field:

- Track reconstruction up to  $|\eta| < 2.47$ ;
- Conversion vertices reconstruction;
- $e/\gamma$  and  $e/\pi^\pm$  separation;
- **Pixel**: (B) 3 layers +(E) 2x3 disks  $\sigma_{r\phi} \sim 10 \mu\text{m}$ ,  $\sigma_z \sim 115 \mu\text{m}$ ;
- **Semi Conductor Tracker**: (B) 4 layers +(E) 2x9 disks  $\sigma_{r\phi} \sim 17 \mu\text{m}$ ,  $\sigma_z \sim 580 \mu\text{m}$ ;
- **Transition Radiation Tracker**: (B) 73 layers +(E) 2x160 layers  $\sigma_z \sim 130 \mu\text{m}$ ;

**LAr lead sampling calorimeter** with an 'accordion' geometry.

- 3 longitudinal layers with cell of  $\Delta\eta \times \Delta\phi$ :
  - 1<sup>st</sup> layer  $(0.003 \div 0.006) \times 0.1$ ;
  - 2<sup>nd</sup> layer  $0.025 \times 0.025$ ;
  - 3<sup>rd</sup> layer  $0.050 \times 0.025$ .
- Presampler for  $|\eta| < 1.8$   $\Delta\eta \times \Delta\phi \sim 0.025 \times 0.1$ .
- Barrel-end-cap crack  $|\eta| = 1.37 \div 1.52$ .
- $\sigma(E)/E = (10-17\%)(\eta)/\sqrt{E(\text{GeV})} \oplus (1.2 \div 1.8\%)$ .









 **ATLAS**  
**EXPERIMENT**

Run Number: 204769, Event Number: 24947130

Date: 2012-06-10 08:17:12 UTC


$$H \rightarrow \gamma\gamma$$






# LHC Performance (2010-2012)

**Spectacular LHC performance  
(rapid increase of data samples)**




**LHC ended pp run at 7+8 TeV  
after delivering more than  $28 \text{ fb}^{-1}$**


**...will come back in 2015 with 13-14 TeV collisions**



**... rapid increase of pile-up conditions**




# Multiple Interactions



**Up to 40 interactions / crossing**  
**(requires enormous efforts to understand the reconstruction of the physics objects...)**

**$Z \rightarrow \mu\mu$  events with  
20 interactions on top**











# Building Blocks






$Z \rightarrow ee, J/\Psi \rightarrow ee \dots$

$eey$  and  $\mu\mu\gamma$




# Detector Material Building Blocks

EM absolute scale







# Building Blocks



Alignment of trackers and muon chambers




Using well-known peaks



# Missing $E_T$ vs PILEUP





# Building Blocks

# TAU ID vs PILEUP



# B-JET TAGGING EFFICIENCY

# JET ENERGY SCALE UNCERTAINTY



# SM Physics


*Selected results on  
jets, photons, W/Z+jets, Top quark,  
Dibosons....*

Just to illustrate the Glory of the SM  
(processes relevant for searches later on...)



# Jet Production



anti- $K_T$  jets with  $R=0.4, 0.6$   
 Jet  $p_T > 20$  GeV,  $|Y| < 4.4$



$$M_{jj} > 260 \text{ GeV}$$

$$y^* = |y^1 - y^2|/2 < 2.5$$

ATLAS-CONF-2012-021



**Inclusive jet and dijet production measured in a wide range of jet  $p_T$ , rapidity and dijet mass.**



**Well described by NLO pQCD predictions**






# Inclusive photons

(cross section for isolated photons)


Phys. Lett. B706 (2011) 150-167  
 Phys. Rev. D83 (2011) 052005  
**ATLAS-CONF-2013-022**



Isolation distribution used to extract the background contributions



**Good agreement with NLO pQCD predictions**  
 (at very low  $E_T^\gamma$  predictions are affected by the limited knowledge of the fragmentation contributions)








# γ+jet


Jet  $p_T > 20$  GeV  
photon  $E_T > 25$  GeV

Measured cross sections with  
 $\eta^\gamma y^{\text{jet}} > 0$  &  $\eta^\gamma y^{\text{jet}} < 0$



Fair agreement with NLO pQCD  
except at very low  $E_T^\gamma$  ( $< 45$  GeV)

difficult region where  
photon purity decreases and  
non-pQCD corrections are sizable






# W+jets



**Very good description of the different Jet  $p_T$  distributions by NLO pQCD and LO ME + PS (ALPGEN)**

**Non trivial test of the ME - PS implementation & matching procedures built inside the MCs**

**→ Input to future MC tunes**





# Z+jets

JHEP07(2013)032




Z ( $\rightarrow \nu\nu$ )+jets irreducible background  
In searches for SUSY, LED, etc....

Z ( $\rightarrow ll$ )+jets fundamental SM measurement...  
→ Very clean samples with no missing  $E_T$







Data well described by NLO pQCD and  
ME + PS (ALPGEN/SHERPA) predictions






# Top




## Simultaneous fit to top mass and JES



# SM Cross Section Summary

Standard Model Total Production Cross Section Measurements


Status: July 2014



# SM Cross Section Summary

## Standard Model Production Cross Section Measurements

Status: July 2014





# Towards the discovery

*Notes on statistics*


*The discovery channels*

# Notes on Statistical Significance

## PHYSICAL REVIEW LETTERS™

Member Subscription Copy  
Library or Other Institutional Use Prohibited Until 2017


Articles published week ending 16 MARCH 2012



Published by  
American Physical Society™

APS  
physics

Volume 108, Number 11



*"This could be the discovery of the century. Depending, of course, on how far down it goes."*

# Likelihood ratio

$$L(\mu, \theta) = f_b \phi_b(m_{\gamma\gamma}) + f_s \phi_s(m_{\gamma\gamma})$$

$$f_s \propto \mu$$



$$n_s = \mu \sigma_s^{\text{visible}}$$

Nuisance parameters

$$\lambda_\mu = \lambda(\mu, \theta) = \frac{L(\mu, \hat{\theta}(\mu))}{L(\hat{\mu}, \hat{\theta})}$$

$$q_\mu = -2 \ln \lambda_\mu$$

$$p_\mu = \int_{q_{\text{obs}}}^{\infty} f(q_\mu | \mu) dq_\mu$$





# Only background ?

Test of “null” hypothesis of no signal




If a real signal appears ...  $p_0 \rightarrow 0$   
(once  $p_0 < 2.87 \times 10^{-7} \rightarrow$  Discovery)

$$\lambda_0 = \lambda(0, \theta) = \frac{L(0, \hat{\theta}(0))}{L(\hat{\mu}, \hat{\theta})}$$

$$q_0 = -2 \ln \lambda_0$$

$$p_0 = \int_{q_{obs}}^{\infty} f(q_0 | 0) dq_0$$



# $CL_s$

(do not exclude your signal...)




$$CL_s = \frac{p_s}{1 - p_b}$$

In the case of very small signals (limited sensitivity) the use of  $p_s$  to exclude signals can lead to false exclusions if the data fluctuates down....

In these cases it is better to use CLs ... which is conservative in the exclusion

If  $CL_s < 0.05 \rightarrow$  excluded at 95% CL



# Only background ?

Test of “null” hypothesis of no signal




If a real signal appears ...  $p_0 \rightarrow 0$   
(once  $p_0 < 2.87 \times 10^{-7} \rightarrow$  Discovery)

$$\lambda_0 = \lambda(0, \theta) = \frac{L(0, \hat{\theta}(0))}{L(\hat{\mu}, \hat{\theta})}$$





$$q_0 = -2 \ln \lambda_0$$

$$p_0 = \int_{q_{obs}}^{\infty} f(q_0 | 0) dq_0$$






# Building up the peaks..



# The Announcement (4<sup>th</sup> July 2012)







(picture: courtesy of A. Hoecker)


# PRESS COVERAGE

after July 4<sup>th</sup> seminars at CERN

CERN black board, Jul. 2012



# Exciting times.... 2011-2012




716  
1

PHYSICS LETTERS B VOL. 716/1 (2012) 1-254


ELSEVIER

Volume 716, Issue 1, 17 September 2012

ISSN 0370-2693





<http://www.elsevier.com/locate/physletb>




# Higgs !!



As in Moriond 2013



A mass peak in the vicinity of 125 GeV



# Higgs !!



A mass peak in the vicinity of 125 GeV







# EPS 2013 High Energy and Particle Physics Prize is awarded to...



**"The ATLAS and CMS collaborations, for the discovery of a Higgs boson, as predicted by the Brout-Englert-Higgs mechanism and to**

**Michel Della Negra, Peter Jenni, and Tejinder Virdee, for their pioneering and outstanding leadership rôles in the making of the ATLAS and CMS experiments"**

**.....it was announced during the LHCP Conference in Barcelona, May 17<sup>th</sup> 2013...**

8<sup>th</sup> Oct. 2013



The Nobel Prize in Physics 2013  
François Englert, Peter Higgs

# The Nobel Prize in Physics 2013



Photo: Pnicolet via  
Wikimedia Commons

**François Englert**



Photo: G-M Greuel via  
Wikimedia Commons

**Peter W. Higgs**

The Nobel Prize in Physics 2013 was awarded jointly to François Englert and Peter W. Higgs *"for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at CERN's Large Hadron Collider"*





*Fundación  
Príncipe de Asturias*

*S.A.R. el Príncipe de Asturias  
Presidente de Honor  
de la Fundación*



► INVESTIGACIÓN  
CIENTÍFICA Y TÉCNICA

Peter Higgs, François  
Englert y el CERN




**“...acuerda por unanimidad conceder el Premio Príncipe de Asturias de Investigación Científica y Técnica 2013 de forma conjunta a los físicos Peter Higgs (Reino Unido) y François Englert (Bélgica) y a la institución internacional CERN, el Laboratorio Europeo de Física de Partículas, por la predicción teórica y detección experimental del Bosón de Higgs.”**

**Oviedo, 29 de Mayo de 2013**

# Higgs at the PDG

Citation: K.A. Olive *et al.* (Particle Data Group), *Chin. Phys. C* **38**, 090001 (2014) (URL: <http://pdg.lbl.gov>)



$$J = 0$$



In the following  $H^0$  refers to the signal that has been discovered in the Higgs searches. Whereas the observed signal is labeled as a spin 0 particle and is called a Higgs Boson, the detailed properties of  $H^0$  and its role in the context of electroweak symmetry breaking need to be further clarified. These issues are addressed by the measurements listed below.

Concerning mass limits and cross section limits that have been obtained in the searches for neutral and charged Higgs bosons, see the sections “Searches for Neutral Higgs Bosons” and “Searches for Charged Higgs Bosons ( $H^\pm$  and  $H^{\pm\pm}$ )”, respectively.

## $H^0$ MASS

A combination of the results from ATLAS and CMS, where a recent unpublished result from CMS is used, yields an average value of  $125.6 \pm 0.3$  GeV, see the review on “Status of Higgs Boson Physics.”

VALUE (GeV)

**$125.7 \pm 0.4$  OUR AVERAGE**

$125.5 \pm 0.2^{+0.5}_{-0.6}$

$125.8 \pm 0.4 \pm 0.4$

DOCUMENT ID

TECN

COMMENT

,2 AAD

13AK ATLS

$pp$ , 7 and 8 TeV

,3 CHATRCHYAN13J

CMS


$pp$ , 7 and 8 TeV

# Higgs at the PDG (as by now)

Citation: K.A. Olive *et al.* (Particle Data Group), *Chin. Phys. C*38, 090001 (2014) (URL: <http://pdg.lbl.gov>)

$H^0$


$J = 0$





# EWK fits vs Higgs



$$m_W^2 \left( 1 - \frac{m_W^2}{m_Z^2} \right) = \frac{\pi\alpha}{\sqrt{2}G_F} (1 + \Delta r)$$



$$\Delta r \sim m_{\text{top}}^2$$

$$\Delta r \sim \ln(m_H)$$

Very remarkable agreement  
(within  $1.3 \sigma$ ) between direct  
 $m_H$  measurement and the indirect  
determination via EWK fits




# EWK fits vs Higgs


Indirect determination of  $M_W$  using measured Higgs mass as input leads to

$$M_W \text{ (indirect)} = 80.359 \pm 0.011 \text{ GeV}$$


→ Better than direct measurement  
(World Average : 15 MeV)



$M_W$  [GeV]



Room for improvement in  $M_W$  measurement  
(sensitive via loops to presence of new physics)



$$J \neq 1$$


$$C = +1$$

$$J^{PC}$$




$0^{++}, 0^{-+}, 2^{++}, 2^{-+} \dots$



# Higgs Spin/Parity





$$CL_s(J_{alt}^P) = \frac{p_0(J_{alt}^P)}{1 - p_0(0^+)}$$






**$J^P = 1^+$  and  $1^-$  rejected at 99.7% CL**  
 **$J^P = 2^+$  rejected at 99.9% CL**  
**→ Evidence for  $J^P = 0^+$**

# Higgs Spin/Parity ( $J^P = 0^+ \text{ vs } 0^-$ )



*Phys. Lett. B 726 (2013), pp. 120*



$$q = \log \frac{L(J^P = 0^+)}{L(J^P = 0^-)}$$

**Data agree with  $0^+$  hypothesis  
 $0^-$  solution excluded at 97.8 % CL**

# Other (Silver) Channels

$$H \rightarrow W^+W^-$$





$$H \rightarrow \tau\tau$$

*WH and ZH ( $H \rightarrow bb$ )*

*ttH*



# Higgs Program in a Glance



Channel categories	ggF 	VBF 	VH 	ttH 
$\gamma\gamma$	✓	✓	✓	✓
ZZ (llll)	✓	✓	✓	✓
WW (llln)	✓	✓	✓	✓
$\tau\tau$	✓	✓	✓	✓
bb	✓	✓	✓	✓
Z $\gamma$	✓	✓		
$\mu\mu$	✓	✓		
Invisible	✓	✓	✓	

*Large Backgrounds*



*Tomorrow.....*

*Rare Decays*

# $H \rightarrow W^+W^- \rightarrow ll\nu\nu$



Limited mass resolution due to the presence of neutrinos




# H → ττ



Analysis in multiple channels with +0/1/2-jets in the final state

2-jet channels optimized for VBF and VH

Considering lepton-lepton, lepton-hadron and hadron-hadron tau decay channels




$$\sigma^{visible} = \mu \sigma_{HIGGS}^{SM}$$






ATLAS Prelim.  
 $m_H = 125 \text{ GeV}$

—  $\sigma(\text{statistical})$   
—  $\sigma(\text{syst. incl. theory})$   
—  $\sigma(\text{theory})$   
Total uncertainty  
 $\pm 1\sigma$  on  $\mu$








# H → ττ





First indication of the Higgs couplings to fermions...






$$qq \rightarrow WH, ZH$$



$$VH \rightarrow bb$$




Intense use of Neural Nets to separate signal from background



At the level of what is expected from the SM




# VH $\rightarrow$ bb

$$qq \rightarrow WH, ZH$$





From Fermilab


Observation at the 3 sigma level












# ttH




## At the edge of the LHC Run I sensitivity...






# Rare Decays



$$\sigma \text{Br}_{\mu\mu} / \sigma^{SM} < 7 @ 95\% \text{ CL}$$



$$\sigma \text{Br}_{Z\gamma} / \sigma^{SM} < 11 @ 95\% \text{ CL}$$



# Tomorrow...

$J^{PC}$

- *Other (Silver) Channels*
- Detailed study on Couplings
- Higgs width
- Invisibly decaying Higgs
- Higgs and Vacuum Stability
- Hierarchy Problem & SUSY
- Search for other Higgs
- What to expect in 2015 -- ?





**End Part I**