


# Non-linear response properties: phenomenology and calculation with TDDFT



David A. Strubbe

Department of Materials Science and Engineering,  
Massachusetts Institute of Technology, Cambridge, MA, USA

Benasque, 9 January 2014


## What is non-linear optics?

Polarizability (linear optics)  $\alpha(-\omega, \omega)$   
absorption, refraction  $\text{Im } \alpha, \text{Re } \alpha$

$$\mu_i(\mathcal{E}) = \mu_{i0} + \alpha_{ij}\mathcal{E}_j + \frac{1}{2}\beta_{ijk}\mathcal{E}_j\mathcal{E}_k + \dots$$


$$\beta(2\omega; -\omega, -\omega)$$

Hyperpolarizability: second-harmonic generation (SHG) *etc.*




Consider only perturbative processes via Taylor expansion of dipole moment.  
Not high-harmonic generation!

# Quantized picture of non-linear optics




absorption

$$\sigma_{\text{abs}} \propto I$$




two-photon absorption

$$\sigma_{\text{TPA}} \propto I^2$$



refraction



second-harmonic generation

$$\sigma_{\text{SHG}} \propto I^2$$

## A zoo of nonlinear optical processes

sum-frequency generation (SFG)	$\beta (\omega_1, \omega_2, -(\omega_1 + \omega_2))$
difference-frequency generation (DFG)	$\beta (\omega_1, \omega_2, -(\omega_1 - \omega_2))$
second-harmonic generation (SHG)	$\beta (\omega, \omega, -2\omega)$
optical rectification	$\beta (\omega, -\omega, 0)$
Pockels (electrooptic) effect	$\beta (\omega, 0, -\omega)$
third-harmonic generation (THG)	$\gamma (\omega, \omega, \omega, -3\omega)$
two-photon absorption	$\text{Im } \gamma (\omega, \omega, -\omega, -\omega)$
four-wave mixing	$\gamma (\omega_1, \omega_2, \omega_3, -(\omega_1 + \omega_2 + \omega_3))$

Energy conservation requires frequency arguments to sum to zero.

## Applications

Characterization in surface science and chemistry (very sensitive)

Optical parametric amplifiers

Pockels cells

Laser pointers

Tunable light sources

Optical logic

coherent anti-Stokes Raman spectroscopy (CARS) (kind of 4-wave mixing)



Two-photon fluorescencing labels in biology

Typically inorganic crystals are used in applications, but organic molecules have the potential to be cheaper and more efficient.

# SHG and SFG at the National Ignition Facility

Lawrence Livermore National Laboratory, Livermore, California

600  $\text{KH}_2\text{PO}_4$  crystals of 400 kg each, ultrapure to avoid absorption.



192 laser beams for inertial confinement fusion

Total power = 500 TW

1053 nm  
1053 nm  
1053 nm

527 nm

351 nm

# The challenges of nonlinear optics: a cautionary tale

VOLUME 7, NUMBER 4

PHYSICAL REVIEW LETTERS

AUGUST 15, 1961

## GENERATION OF OPTICAL HARMONICS\*

P. A. Franken, A. E. Hill, C. W. Peters, and G. Weinreich

The Harrison M. Randall Laboratory of Physics, The University of Michigan, Ann Arbor, Michigan

(Received July 21, 1961)




FIG. 1. A direct reproduction of the first plate in which there was an indication of second harmonic. The wavelength scale is in units of 100 A. The arrow at 3472 A indicates the small but dense image produced by the second harmonic. The image of the primary beam at 6943 A is very large due to halation.

The original experimental report of SHG (quartz, 694 nm)

Lesson: check your proofs!

## Symmetry properties of nonlinear susceptibility tensors

Inversion symmetry: even orders are zero ( $\mu$ ,  $\beta$ ,  $\delta$ , etc.).

$$\mu^{(2)} = \frac{1}{2}\beta\mathcal{E}^2$$

$$\text{Apply inversion: } -\mu^{(2)} = \frac{1}{2}\beta(-\mathcal{E})^2 = \frac{1}{2}\beta\mathcal{E}^2 = \mu^{(2)}$$

Therefore  $\beta = 0$  Hence surface-sensitive if bulk is centrosymmetric.

Permutation symmetry:

$$\beta_{ijk}(\omega_1, \omega_2, \omega_3) = \frac{\partial^3 E}{\partial \mathcal{E}_{i,\omega_1} \partial \mathcal{E}_{j,\omega_2} \partial \mathcal{E}_{k,\omega_3}} = \beta_{jik}(\omega_2, \omega_1, \omega_3)$$

$$\beta_{ijk}(\omega_1, \omega_2, \omega_3) \neq \beta_{jik}(\omega_1, \omega_2, \omega_3) \quad \text{“Kleinman symmetry,”}$$


Only for  $\omega \approx 0$

Also Kramers-Kronig relations.



# Symmetry properties of nonlinear susceptibility tensors

Spatial symmetries: e.g. chloroform ( $\text{CHCl}_3$ )



## Character table for point group $C_{3v}$

$C_{3v}$	E	$2C_3$ (z)	$3\sigma_v$	linear functions, rotations	quadratic functions	cubic functions
$A_1$	+1	+1	+1	z	$x^2+y^2, z^2$	$z^3, x(x^2-3y^2), z(x^2+y^2)$
$A_2$	+1	+1	-1	$R_z$	-	$y(3x^2-y^2)$
E	+2	-1	0	(x, y) ( $R_x,$ $R_y$ )	$(x^2-y^2, xy)$ (xz, yz)	$(xz^2, yz^2)$ [xyz, $z(x^2-y^2)$ ] [ $x(x^2+y^2),$ $y(x^2+y^2)$ ]

$$\alpha_{xy} = \frac{\partial^2 E}{\partial x \partial y} \sim xy \quad \begin{matrix} \uparrow \\ \mu_j \end{matrix} \quad \begin{matrix} \uparrow \\ \alpha_{ij} \end{matrix} \quad \begin{matrix} \uparrow \\ \beta_{ijk} \end{matrix}$$

Zincblende structure (e.g. GaAs) has only  $\chi_{xyz}^{(2)} \neq 0$

# Conventions and units

Many conventions for pre-factors! Multiple papers on just conventions...

TABLE I. Expressions for the total dipole moment measured in a dc-electric field induced second harmonic generation experiment

Label	Convention	$[\mu_{\text{ind}}(2\omega)]/(F_0 F_\omega^2 \cos 2\omega t)$	Zero freq. convergence <sup>a</sup>	
			$\rightarrow \beta_0$	$\rightarrow \gamma_0$
$AB \equiv T$	Taylor series	$\frac{1}{4} \left[ \tilde{\gamma}^T(-2\omega; \omega, \omega, 0) + \frac{\mu\beta_z^T(-2\omega; \omega, \omega)}{5kT} \right]$	+	+
$B$	Perturbation series	$\frac{3}{2} \left[ \tilde{\gamma}^B(-2\omega; \omega, \omega, 0) + \frac{\mu\beta_z^B(-2\omega; \omega, \omega)}{15kT} \right]$	+	+
$B^*$	EFISH only	$\frac{3}{2} \left[ \tilde{\gamma}^{B^*}(-2\omega; \omega, \omega, 0) + \frac{\mu\beta_z^{B^*}(-2\omega; \omega, \omega)}{5kT} \right]$	-	+
$A$	Not used	$\frac{1}{6} \left[ \tilde{\gamma}^A(-2\omega; \omega, \omega, 0) + \frac{3\mu\beta_z^A(-2\omega; \omega, \omega)}{5kT} \right]$	-	-
$X$	Phenomenological	$\left[ \tilde{\gamma}^X(-2\omega; \omega, \omega, 0) + \frac{\mu\beta_z^X(-2\omega; \omega, \omega)}{5kT} \right]$	-	-

<sup>a</sup>Columns three and four indicate whether the convention does (+) or does not (-) converge to  $\beta_0$  or  $\gamma_0$  as  $\omega \rightarrow 0$ .

Commonly used units: au, esu

$$\begin{aligned}
 &1 \text{ au of } \beta \\
 &= 3.206361 \times 10^{-53} \text{ C}^3 \text{ m}^3/\text{J}^2 \text{ (SI)} \\
 &= 8.6392 \times 10^{-33} \text{ cm}^4 \text{ statvolt}^{-1} \text{ (esu of } \beta)
 \end{aligned}$$

A. Willetts, J. E. Rice, D. M. Burland, and D. P. Shelton, *J. Chem. Phys.* **97**, 7590 (1992)

## Nonlinear optics in solids

Susceptibility tensors for solids  $\chi_{ij}^{(1)} = \alpha_{ij}/V$   $\chi_{ij}^{(2)} = \beta_{ij}/V$


Now can have  $q$ -dependence as well.  $\chi_{ijk}^{(2)}(\vec{q}_1, \vec{q}_2, \vec{q}_3, \omega_1, \omega_2, \omega_3)$

Phase-matching condition for constructive interference:

$$\vec{k}_1 + \vec{k}_2 + \vec{k}_3 = 0 \quad (\text{NOT momentum conservation})$$
$$= \frac{n_i(\omega_1)\omega_1}{c} + \frac{n_j(\omega_2)\omega_2}{c} + \frac{n_k(\omega_3)\omega_3}{c}$$

Adjust angle of incidence to satisfy and get significant conversion.

Molecules vs solids: orders of magnitude.



$\beta \sim 10000$  au

KTiOPO<sub>4</sub>  $\chi^{(2)} \sim 10^{-8}$  esu

CHCl<sub>3</sub>  $\beta \sim 1$  au,  $\chi^{(2)} \sim 10^{-8}$  esu

## Other perturbations

Ionic displacement and strain (Raman tensors, Grüneisen parameters, phonon anharmonicity, pyroelectric tensor, piezoelectric tensor, second-order elastic coefficients, ...)

S. Baroni, S. de Gironcoli, A. Dal Corso, and P. Gianozzi, *Rev. Mod. Phys.* **73**, 515 (2001)

Magnetic contributions to nonlinear optical processes.

But usually electric-dipole approximation is sufficient!

Representation of electric field in finite and periodic systems:

$$r \rightarrow -i \frac{\partial}{\partial k} \quad V_{\mathcal{E}} = -i \frac{\partial}{\partial k} \quad \text{Quantum theory of polarization}$$

A Dal Corso, F Mauri, and A Rubio, *Phys. Rev. B* **53**, 15638 (1996)

## Solution measurements of hyperpolarizability

Measurements are usually in solution for molecules. Solvent effects can be strong and complicate comparison between experiment and theory.

(In theory, handle via polarizable continuum models or explicit solvent in small clusters or periodic system.)

(Time-averaged) inversion symmetry of solution makes ordinary measurement give zero.


Electric-field-induced second-harmonic generation (EFISH) is coherent third-order process, based on field lining up molecules.

$$\chi^{(3)}(-2\omega; \omega, \omega, 0) = n \left[ \gamma(-2\omega; \omega, \omega, 0) + \frac{\mu}{3kT} \beta_{\parallel}(-2\omega; \omega, \omega) \right]$$

What is measured directly and often reported:  $\mu\beta_{\parallel}$

Hyper-Rayleigh scattering (HRS) is incoherent second-order process, based on orientational fluctuations.

$$I_{\text{HRS}} \propto \langle \beta_{xyz} \beta_{uvw} \rangle$$



## Solution measurements of hyperpolarizability

Consider projection along dipole moment.

$$\beta_{\text{EFISH}}^i = \frac{1}{5} \sum_j (\beta_{ijj} + \beta_{jij} + \beta_{jji})$$

Vertical-vertical and horizontal-vertical polarizations for experiment.

$$[\beta_{\text{HRS}}^{\text{VV}}]^2 = \frac{1}{7} \sum_i \beta_{iii}^2 + \frac{6}{35} \sum_{i \neq j} \beta_{iii} \beta_{ijj} + \frac{9}{35} \sum_{i \neq j} \beta_{ijj}^2 + \frac{6}{35} \sum_{\text{cyclic}} \beta_{ijj} \beta_{jkk} + \frac{12}{35} \beta_{ijk}^2$$

$$[\beta_{\text{HRS}}^{\text{HV}}]^2 = \frac{1}{35} \sum_i \beta_{iii}^2 - \frac{2}{105} \sum_{i \neq j} \beta_{iii} \beta_{ijj} + \frac{11}{105} \sum_{i \neq j} \beta_{ijj}^2 - \frac{2}{105} \sum_{\text{cyclic}} \beta_{ijj} \beta_{jkk} + \frac{8}{35} \beta_{ijk}^2$$

## Vibrational / rotational contributions

Many measurements are at  $\lambda = 1064$  nm. For organic molecules, typically:


- above vibrational frequencies, so can neglect vibrations
- below electronic resonances so little dispersion

Rotational and vibrational contributions can be estimated from simple harmonic models, for low-frequency response (e.g. THz).

$Z^*$  = Born effective charge

$$F = \mathcal{E}Z^* - kx = 0$$

$$\beta^{\text{vib}} = \frac{Z}{k} \left( 2 \frac{\partial Z}{\partial \mathcal{E}} - \frac{Z}{k} \right)$$



$$\beta^{\text{rot}} = \frac{3\mu}{kT} \alpha$$

D. Bishop, *Rev. Mod. Phys.* **62**, 343 (1990)

E. Roman, J. R. Yates, M. Veithen, D. Vanderbilt, and I. Souza, *Phys. Rev. B* **74**, 245204 (2006)



## Local-field factors for solvent

$$\mathcal{E}_{\text{screened}} = \mathcal{E}_{\text{external}} / \epsilon$$

Clausius-Mossotti relations,  
relate bulk to molecular susceptibilities.

$$\mathcal{E}_{\text{local}} = f \mathcal{E}_{\text{screened}}$$

$$\chi^{(2)} = \rho f_{\omega}^2 f_{2\omega} \beta$$



Also generalizations for ellipsoidal cavity, polar solvents.







## Theoretical methods for nonlinear response

1. Finite differences (static)
2. Explicit time-propagation
3. Sternheimer equation ( $2n+1$  Theorem)
4. Sum over states (Casida equation)
5. Dyson equation

# Finite differences

Apply static fields and calculate dipole moment.  
No need for special capabilities in code.  
Probably most common method used.  
Hope static and IR are similar!



F. Vila, D.A. Strubbe, Y. Takimoto, X. Andrade, A. Rubio, S. G. Louie, and J. J. Rehr, *J. Chem. Phys.* **133**, 034111 (2010)

## Convergence is more demanding for nonlinear response


Basis Set	$\mu_z$	$\alpha_{yy}$	$\alpha_{zz}$	$\beta_{yyy}$	$\beta_{yyz}$	$\beta_{zzz}$	$\bar{\alpha}$	$\beta_{\parallel}$	$\beta_{\text{HRS}}^{\text{VV}}$
GTO 5Z <sub>sa</sub> (aug-cc-pV5Z)	0.404	65.70	46.79	27.35	-15.31	22.27	59.40	-5.01	16.90
NBS 5Z4P <sub>e8</sub> (SIESTA)	0.398	65.45	46.28	24.54	-14.90	21.37	59.06	-5.07	15.68
RS 1r (h=0.25, r=22)	0.399	66.02	47.00	27.12	-16.36	26.94	59.68	-3.47	17.44
RS fd	"	66.46	47.07	24.22	-15.66	25.50	60.00	-3.52	16.14
RS 1064 nm	"	66.69	47.34	30.35	-18.95	31.56	60.24	-4.01	19.91
<b>Expt.</b>	$0.409 \pm 0.008$	$61 \pm 5$	$45 \pm 3$				$56 \pm 4$	$1 \pm 4$	

F. Vila, D.A. Strubbe, Y. Takimoto, X. Andrade, A. Rubio, S. G. Louie, and J. J. Rehr, *J. Chem. Phys.* **133**, 034111 (2010)

Comparison of Gaussian-type orbitals, SIESTA numerical basis sets, real-space grid: need five-zeta basis sets with diffuse functions, or very long-range real-space grid.




Response extends far from molecule ( $> 15 a_0$ )

Compare: only  $12 a_0$  converges ground state  
all atoms contained within  $3 a_0$  radius



## Explicit time-propagation



Cannot use “kick” for all frequencies at once (as for linear optics).  
Must calculate separately for each frequency combination.  
Scaling is like one ground-state calculation per time step.  
Incident laser pulses (duration  $\sim$  imaginary broadening).



p-nitroaniline

$$P_{ijk}^{(2)}(\omega) = \frac{D^{(2)}}{2\pi} \chi_{ijk}^{(2)}(-\omega; \omega_1, \omega_2) F_{\omega_1} F_{\omega_2}$$

# Explicit time-propagation



## Sternheimer equation

*aka* density-functional perturbation theory or coupled perturbed Kohn-Sham

Calculate variation of wavefunctions in linear response.

No need for unoccupied states.

One frequency at a time (but can use previous freq as starting point).

$$\left( H^{(0)} - \epsilon^{(0)} \pm \omega_{\alpha} \right) P'_n \psi_{\alpha\pm}^{(1)} = -P'_n H_{\alpha\pm}^{(1)} \psi^{(0)}$$

$$H_{\alpha\pm}^{(1)} = V_{\alpha\pm}^{(1)} + V_H \left[ n_{\alpha\pm}^{(1)} \right] + \int f_{xc} [n] n_{\alpha\pm}^{(1)}(r) d^3r \quad \text{SCF cycle}$$

$$n_{\pm}^{(1)} = \sum_n^{\text{occ}} \left( \psi_{n\pm}^{(1)} [\psi_n^{(0)}]^* + \psi_n^{(0)} [\psi_{n\mp}^{(1)}]^* \right)$$

$$P'_n = 1 - \left| \psi_n^{(0)} \right\rangle \left\langle \psi_n^{(0)} \right| \quad \left\langle \psi^{(0)} \left| \psi_{\alpha\pm}^{(1)} \right\rangle = 0$$

Need small imaginary broadening  $i\eta$  near resonances for numerical stability.

## Sternheimer equation: $2n+1$ Theorem

Solving  $n^{\text{th}}$ -order perturbation theory gives  $2n+1$  derivative of total energy.

$$\chi^{(2n)} = \frac{\partial^{(2n+1)} E}{\partial \lambda^{(2n+1)}} \quad F = \frac{\partial E}{\partial R} = \left\langle \psi \left| \frac{\partial H}{\partial R} \right| \psi \right\rangle$$

$n = 0$ : Hellman-Feynman Theorem. No wavefunction derivatives at all.

$n = 1$ : Linear variation of wavefunctions gives quadratic response.


$$\begin{aligned} \beta_{ijk}(-\omega_1; \omega_2, \omega_3) = & -4 \sum_P \sum_{\zeta=\pm 1} \left[ \sum_m^{\text{occ}} \left\langle \psi_{mi}^{(1)}(-\zeta\omega_1) \left| H_j^{(1)}(\zeta\omega_2) \right| \psi_{mk}^{(1)}(\zeta\omega_3) \right\rangle \right. \\ & - \sum_{mn}^{\text{occ}} \left\langle \psi_m^{(0)} \left| H_j^{(1)}(\zeta\omega_2) \right| \psi_m^{(0)} \right\rangle \left\langle \psi_{mi}^{(1)}(-\zeta\omega_1) \left| \psi_{mk}^{(1)}(\zeta\omega_3) \right\rangle \right. \\ & \left. \left. - \frac{2}{3} \int d^3r \int d^3r' \int d^3r'' K_{xc}(r, r', r'') n_i^{(1)}(r, \omega_1) n_j^{(1)}(r', \omega_2) n_k^{(1)}(r'', \omega_3) \right] \right. \end{aligned}$$

X. Gonze and J.-P. Vigneron, *Phys. Rev. B* **39**, 13120 (1989)

Quantum ESPRESSO and Octopus use equations on previous slide.



ABINIT formulation: find  $\beta$  by minimizing with respect to  $\psi^{(1)}$  (variational).

# Sternheimer equation: examples from Octopus code



CO linear spectrum

## H<sub>2</sub>O optical rectification



CO SHG



X Andrade, S Botti, MAL Marques, and A Rubio, *J. Chem. Phys.* **126**, 184106 (2007)



# Sternheimer equation: examples from Octopus code

Efficient scaling with system size: same as ground state.

e.g. comparison of TDLDA vs. CIS with Gaussian basis for vibrating N@C<sub>60</sub>



## Sum over states

Applicable to any method giving excited-state energies and matrix elements, in particular Casida equation for TDDFT.

(Also other theories such as configuration interaction, coupled cluster, Bethe-Salpeter equation, etc.)

Most commonly used as RPA though: just use Kohn-Sham energies and matrix elements.

## Sum over states

Arbitrary (or physical but unknown) imaginary broadenings  $\Gamma$ .

Convergence in two sums on states is difficult!

Applied occasionally to two-photon absorption in TDDFT.

$$\begin{aligned}\chi_{ij}^{(1)}(\omega) &= \frac{P_i^{(1)}(\omega)}{E_j(\omega)} \\ &= N \frac{e^2}{\hbar} \sum_{gn} \left[ \frac{(r_i)_{ng}(r_j)_{gn}}{\omega + \omega_{ng} + i\Gamma_{ng}} - \frac{(r_j)_{ng}(r_i)_{gn}}{\omega - \omega_{ng} + i\Gamma_{ng}} \right] \rho_g^{(0)},\end{aligned}$$

$$\begin{aligned}\chi_{ijk}^{(2)}(\omega = \omega_1 + \omega_2) &= \frac{P_i^{(2)}(\omega)}{E_j(\omega_1)E_k(\omega_2)} \\ &= -N \frac{e^3}{\hbar^2} \sum_{g,n,n'} \left[ \frac{(r_i)_{gn}(r_j)_{nn'}(r_k)_{n'g}}{(\omega - \omega_{ng} + i\Gamma_{ng})(\omega_2 - \omega_{n'g} + i\Gamma_{n'g})} \right. \\ &\quad + \frac{(r_i)_{gn}(r_k)_{nn'}(r_j)_{n'g}}{(\omega - \omega_{ng} + i\Gamma_{ng})(\omega_1 - \omega_{n'g} + i\Gamma_{n'g})} \\ &\quad + \frac{(r_k)_{gn'}(r_j)_{n'n}(r_i)_{ng}}{(\omega + \omega_{ng} + i\Gamma_{ng})(\omega_2 + \omega_{n'g} + i\Gamma_{n'g})} \\ &\quad + \frac{(r_j)_{gn'}(r_k)_{n'n}(r_i)_{ng}}{(\omega + \omega_{ng} + i\Gamma_{ng})(\omega_1 + \omega_{n'g} + i\Gamma_{n'g})} \\ &\quad - \frac{(r_j)_{ng}(r_i)_{n'n}(r_k)_{gn'}}{(\omega - \omega_{nn'} + i\Gamma_{nn'})} \left( \frac{1}{\omega_2 + \omega_{n'g} + i\Gamma_{n'g}} + \frac{1}{\omega_1 - \omega_{ng} + i\Gamma_{ng}} \right) \\ &\quad \left. - \frac{(r_k)_{ng}(r_i)_{n'n}(r_j)_{gn'}}{(\omega - \omega_{nn'} + i\Gamma_{nn'})} \left( \frac{1}{\omega_2 - \omega_{ng} + i\Gamma_{ng}} + \frac{1}{\omega_1 + \omega_{n'g} + i\Gamma_{n'g}} \right) \right] \rho_g^{(0)}.\end{aligned}\tag{2.17}$$

## Dyson-like equation

Efficient scheme for solids (for linear and non-linear optics):  $k$ -points.  
Need  $q \rightarrow 0$  limit, equivalent to  $k \cdot p$  perturbation theory.

$$\begin{aligned}
 & [1 - \chi_0^{(1)}(\omega_1 + \omega_2) f_{uxc}(\omega_1 + \omega_2)] \chi_{\rho\rho\rho}^{(2)}(\omega_1, \omega_2) \\
 &= \chi_0^{(2)}(\omega_1, \omega_2) [1 + f_{uxc}(\omega_1) \chi^{(1)}(\omega_1)] \\
 &\quad \times [1 + f_{uxc}(\omega_2) \chi^{(1)}(\omega_2)] + \chi_0^{(1)}(\omega_1 + \omega_2) g_{xc}(\omega_1 + \omega_2) \\
 &\quad \times \chi^{(1)}(\omega_1) \chi^{(1)}(\omega_2),
 \end{aligned}$$

Complicated macroscopic/microscopic relations...

G. Senatore and K. R. Subbaswamy, *Phys. Rev. A*  
**35**, 2440 (1987)

E. Luppi, H. Hübener, and V. Véniard, *J. Chem. Phys.*  
**132**, 241104 (2010)




FIG. 1. Experimental spectrum of GaAs from Bergfeld and Daum (Ref. 23) (circles). Calculated  $|\chi_{xyz}^{(2)}|$ : IPA (solid line); including crystal local-field effects (dashed line); including excitonic effects through the long-range kernel with theoretical (dotted-dashed line) and experimental (Ref. 36) (thick solid line)  $\epsilon_M^L$ ; theoretical calculation from Leitsmann *et al.* (Ref. 17) where the excitons are included within BSE framework (diamond). It should be noted that no scaling or fitting has been done for any of the curves.

## Dyson-like equation

Must converge with respect to unoccupied states in a triple sum.

$$\begin{aligned}
 \chi_{0,G,G_1,G_2}^{(2)}(2\mathbf{q},\mathbf{q},\mathbf{q},\omega) = & \frac{2}{V} \sum_{n,n',n'',\mathbf{k}} \frac{\langle \phi_{n,\mathbf{k}} | e^{-i(2\mathbf{q}+\mathbf{G})\mathbf{r}} | \phi_{n',\mathbf{k}+2\mathbf{q}} \rangle}{(E_{n,\mathbf{k}} - E_{n',\mathbf{k}+2\mathbf{q}} + 2\omega + 2i\eta)} \left[ (f_{n,\mathbf{k}} - f_{n'',\mathbf{k}+q}) \frac{\langle \phi_{n',\mathbf{k}+2\mathbf{q}} | e^{i(\mathbf{q}+\mathbf{G}_1)\mathbf{r}_1} | \phi_{n'',\mathbf{k}+q} \rangle \langle \phi_{n'',\mathbf{k}+q} | e^{i(\mathbf{q}+\mathbf{G}_2)\mathbf{r}_2} | \phi_{n,\mathbf{k}} \rangle}{(E_{n,\mathbf{k}} - E_{n'',\mathbf{k}+q} + \omega + i\eta)} \right. \\
 & + (f_{n,\mathbf{k}} - f_{n'',\mathbf{k}+q}) \frac{\langle \phi_{n',\mathbf{k}+2\mathbf{q}} | e^{i(\mathbf{q}+\mathbf{G}_2)\mathbf{r}_2} | \phi_{n'',\mathbf{k}+q} \rangle \langle \phi_{n'',\mathbf{k}+q} | e^{i(\mathbf{q}+\mathbf{G}_1)\mathbf{r}_1} | \phi_{n,\mathbf{k}} \rangle}{(E_{n,\mathbf{k}} - E_{n'',\mathbf{k}+q} + \omega + i\eta)} \\
 & + (f_{n',\mathbf{k}+2\mathbf{q}} - f_{n'',\mathbf{k}+q}) \frac{\langle \phi_{n',\mathbf{k}+2\mathbf{q}} | e^{i(\mathbf{q}+\mathbf{G}_2)\mathbf{r}_2} | \phi_{n'',\mathbf{k}+q} \rangle \langle \phi_{n'',\mathbf{k}+q} | e^{i(\mathbf{q}+\mathbf{G}_1)\mathbf{r}_1} | \phi_{n,\mathbf{k}} \rangle}{(E_{n'',\mathbf{k}+q} - E_{n',\mathbf{k}+2\mathbf{q}} + \omega + i\eta)} \\
 & \left. + (f_{n',\mathbf{k}+2\mathbf{q}} - f_{n'',\mathbf{k}+q}) \frac{\langle \phi_{n',\mathbf{k}+2\mathbf{q}} | e^{i(\mathbf{q}+\mathbf{G}_1)\mathbf{r}_1} | \phi_{n'',\mathbf{k}+q} \rangle \langle \phi_{n'',\mathbf{k}+q} | e^{i(\mathbf{q}+\mathbf{G}_2)\mathbf{r}_2} | \phi_{n,\mathbf{k}} \rangle}{(E_{n'',\mathbf{k}+q} - E_{n',\mathbf{k}+2\mathbf{q}} + \omega + i\eta)} \right], \quad (38)
 \end{aligned}$$

## Some references on nonlinear response

DA Strubbe, L Lehtovaara, A Rubio, MAL Marques, and SG Louie, “Response functions in TDDFT: concepts and implementation,” in *Fundamentals of Time-dependent density-functional theory* (ch. 7) and references therein.

Yuen-Ron Shen, *The Principles of Nonlinear Optics* (Wiley 1984 / 2002).

P. N. Butcher and D. Cotter, *The Elements of Nonlinear Optics* (Cambridge University Press, 1990).