

# Bethe-Salpeter equation: electron-hole excitations and optical spectra

Ilya Tokatly

*European Theoretical Spectroscopy Facility (ETSF)*

*NanoBio Spectroscopy Group - UPV/EHU San Sebastián - Spain*

*IKERBASQUE, Basque Foundation for Science - Bilbao - Spain*

[ilya.tokatly@ehu.es](mailto:ilya.tokatly@ehu.es)

TDDFT school - Benasque 2018


# Outline

- 1 Optics and two-particle dynamics: Why BSE?
- 2 The Bethe-Salpeter equation: Pictorial derivation
- 3 Macroscopic response and the Bethe-Salpeter equation
- 4 The Bethe-Salpeter equation in practice


# Outline

- 1 Optics and two-particle dynamics: Why BSE?
- 2 The Bethe-Salpeter equation: Pictorial derivation
- 3 Macroscopic response and the Bethe-Salpeter equation
- 4 The Bethe-Salpeter equation in practice

# Optical absorption: Experiment and Phenomenology



Silicon  
Optical Absorption



Exp. at 30 K from: P. Lautenschlager *et al.*,  
Phys. Rev. B **36**, 4821 (1987).

Light is absorbed:  $I = I_0 e^{-\alpha(\omega)x}$

Classical electrodynamics

$$E = E_0 e^{-i(\omega t - qx)}, \quad q^2 = \frac{\omega^2}{c^2} \epsilon_M(\omega)$$

$$\epsilon_M(\omega) = \epsilon'_M(\omega) + i\epsilon''_M(\omega)$$

$$q \approx \frac{\omega}{c} \sqrt{\epsilon'_M} + i \frac{\omega}{2c\sqrt{\epsilon'_M}} \epsilon''_M$$

$$\sqrt{\epsilon'_M} = n_r - \text{index of refraction}$$


$$I \sim |E|^2 = |E_0|^2 e^{-\alpha(\omega)x}$$

$$\alpha(\omega) = \frac{\omega}{cn_r} \epsilon''_M(\omega)$$

$$\epsilon''_M(\omega) \sim \text{absorption rate}$$


# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair



# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair



# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair

**Representation by Feynman diagrams:**



- photon creates an e-h pair
- the pair propagates freely
- it recombines and recreates a photon

# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair

**Representation by Feynman diagrams:**



- photon creates an e-h pair
- the pair propagates freely
- it recombines and recreates a photon



# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair

**Representation by Feynman diagrams:**



- photon creates an e-h pair
- the pair propagates freely
- it recombines and recreates a photon

# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair

**Representation by Feynman diagrams:**



- photon creates an e-h pair
- the pair propagates freely
- it recombines and recreates a photon

# Optical absorption: Microscopic picture

Elementary process of absorption: Photon creates a single e-h pair

**Representation by Feynman diagrams:**



- photon creates an e-h pair
- the pair propagates freely
- it recombines and recreates a photon

Absorption rate is given by an imaginary part of the polarization loop



$$W = \frac{2\pi}{\hbar} \sum_{i,j} |\langle \varphi_i | \mathbf{e} \cdot \hat{\mathbf{v}} | \varphi_j \rangle|^2 \delta(\varepsilon_j - \varepsilon_i - \hbar\omega) \sim \text{Im}\epsilon(\omega)$$

# Absorption by independent Kohn-Sham particles



Independent transitions:

$$\epsilon''(\omega) = \frac{8\pi^2}{\omega^2} \sum_{ij} |\langle \varphi_j | \mathbf{e} \cdot \hat{\mathbf{v}} | \varphi_i \rangle|^2 \delta(\epsilon_j - \epsilon_i - \omega)$$





# Absorption by independent Kohn-Sham particles



Independent transitions:


$$\epsilon''(\omega) = \frac{8\pi^2}{\omega^2} \sum_{ij} |\langle \varphi_j | \mathbf{e} \cdot \hat{\mathbf{v}} | \varphi_i \rangle|^2 \delta(\epsilon_j - \epsilon_i - \omega)$$



Particles are interacting!

# Interaction effects: self-energy corrections


## 1st class of interaction corrections:



Created electron and hole interact with other particles in the system,  
but do not touch each other


# Interaction effects: self-energy corrections

## 1st class of interaction corrections:



Created electron and hole interact with other particles in the system, but do not touch each other

## Absorption by “dressed” particles



Bare propagator  $G_0$  is replaced by the full propagator  
 $G = G_0 + G_0 \Sigma G$

$$[\omega - \hat{h}_0(\mathbf{r})]G(\mathbf{r}, \mathbf{r}', \omega) + \int d\mathbf{r}_1 \Sigma(\mathbf{r}, \mathbf{r}_1, \omega)G(\mathbf{r}_1, \mathbf{r}', \omega) = \delta(\mathbf{r} - \mathbf{r}')$$

# Self-energy corrections

## Perturbative GW corrections

$$\hat{h}_0(\mathbf{r})\varphi_i(\mathbf{r}) + V_{xc}(\mathbf{r})\varphi_i(\mathbf{r}) = \epsilon_i\varphi_i(\mathbf{r})$$

$$\hat{h}_0(\mathbf{r})\phi_i(\mathbf{r}) + \int d\mathbf{r}' \Sigma(\mathbf{r}, \mathbf{r}', \omega = E_i) \phi_i(\mathbf{r}') = E_i \phi_i(\mathbf{r})$$

First-order perturbative corrections with  $\Sigma = GW$ :

$$E_i - \epsilon_i = \langle \varphi_i | \Sigma - V_{xc} | \varphi_i \rangle$$

Hybersten and Louie, PRB **34** (1986);  
Godby, Schlüter and Sham, PRB **37** (1988)





# Optical absorption: Independent quasiparticles




Independent transitions:

$$\epsilon''(\omega) = \frac{8\pi^2}{\omega^2} \sum_{ij} |\langle \varphi_j | \mathbf{e} \cdot \hat{\mathbf{v}} | \varphi_i \rangle|^2 \delta(E_j - E_i - \omega)$$



# Interaction effects: vertex (excitonic) corrections


## 2nd class of interaction corrections:



includes all direct and indirect interactions between electron and hole created by a photon

# Interaction effects: vertex (excitonic) corrections

## 2nd class of interaction corrections:



includes all direct and indirect interactions between electron and hole created by a photon


Summing up all such interaction processes we get:



Empty polarization loop is replaced by the full two-particle propagator  
 $L(\mathbf{r}_1 t_1; \mathbf{r}_2 t_2; \mathbf{r}_3 t_3; \mathbf{r}_4 t_4) = L(1234)$  with joined ends

# Interaction effects: vertex (excitonic) corrections

## 2nd class of interaction corrections:



includes all direct and indirect interactions between electron and hole created by a photon

Summing up all such interaction processes we get:




Empty polarization loop is replaced by the full two-particle propagator

$$L(\mathbf{r}_1 t_1; \mathbf{r}_2 t_2; \mathbf{r}_3 t_3; \mathbf{r}_4 t_4) = L(1234) \text{ with joined ends}$$

**Equation for  $L(1234)$  is the Bethe-Salpeter equation!**


# Absorption



Neutral excitations  $\rightarrow$  poles of two-particle Green's function  $L$

Excitonic effects = electron - hole interaction


# Absorption



Neutral excitations  $\rightarrow$  poles of two-particle Green's function  $L$

Excitonic effects = electron - hole interaction

# Absorption



Neutral excitations  $\rightarrow$  poles of two-particle Green's function  $L$

Excitonic effects = electron - hole interaction


# Outline

- 1 Optics and two-particle dynamics: Why BSE?
- 2 The Bethe-Salpeter equation: Pictorial derivation**
- 3 Macroscopic response and the Bethe-Salpeter equation
- 4 The Bethe-Salpeter equation in practice



# Derivation of the Bethe-Salpeter equation (1)

Propagator of e-h pair in a many-body system:



- Solid lines stand for bare one-particle Green's functions

$$G_0(12) = G_0(\mathbf{r}_1, \mathbf{r}_2, t_1 - t_2)$$


- Wiggled lines correspond to the interaction (Coulomb) potential

$$v(12) = v(\mathbf{r}_1 - \mathbf{r}_2)\delta(t_1 - t_2) = \frac{e^2}{|\mathbf{r}_1 - \mathbf{r}_2|}\delta(t_1 - t_2)$$


- Integration over space-time coordinates of all intermediate points in each graph is assumed

# Derivation of the Bethe-Salpeter equation (1)


Propagator of e-h pair in a many-body system:




**1st step: Dressing one-particle propagators**



Self-energy  $\Sigma$  is a sum of all 1-particle irreducible diagrams




Full 1-particle Green's function satisfies the Dyson equation




# Derivation of the Bethe-Salpeter equation (2)

Propagation of dressed interacting electron and hole:




## 2nd step: Classification of scattering processes

At this stage we identify two-particle irreducible blocks



where  $\gamma(1234)$  of the electron-hole scattering amplitude



# Derivation of the Bethe-Salpeter equation (3)

**Final step: Summation of a geometric series**

$$L = \text{---} + \Theta + \Theta L + \dots$$


**The result is the Bethe-Salpeter equation**

$$L = \text{---} + \Theta L$$


$$\Theta = \text{---} - \Gamma$$

# Derivation of the Bethe-Salpeter equation (3)

**Final step: Summation of a geometric series**



**The result is the Bethe-Salpeter equation**



**Analytic form of the Bethe-Salpeter equation ( $j = \{\mathbf{r}_j, t_j\}$ )**

$$L(1234) = L_0(1234) + \int L_0(1256)[v(57)\delta(56)\delta(78) - \gamma(5678)]L(7834)d5d6d7d8$$

# Closed set of equations in a diagrammatic form

The diagrammatic Dyson equation for the  $L$ -vertex is shown as:

$$\boxed{L} = \text{two parallel lines} + \text{self-energy loop} \cdot \boxed{L} - \text{vertex correction } \gamma \cdot \boxed{L}$$

- 1-particle Green's function  $G(12)$  satisfies the Dyson equation

The Dyson equation for the 1-particle Green's function is shown as:

$$\text{two parallel lines} = \text{single line} + \text{single line} \cdot \Sigma \cdot \text{two parallel lines}$$

- $\Sigma(12)$  is a sum of all 1-particle irreducible diagrams

The expansion of the self-energy vertex  $\Sigma$  is shown as:

$$\Sigma = \text{cloud} + \text{cloud with bubble} + \text{cloud with bubble and loop} + \dots$$

- $\gamma(1234)$  – sum of all e-h and interaction irreducible diagrams

The expansion of the vertex correction  $\gamma$  is shown as:

$$\gamma = \text{wavy line} + \text{wavy line with bubble} + \text{wavy line with bubble and cloud} + \text{wavy line with bubble and cloud and wavy line} + \dots = \delta\Sigma/\delta G$$

# Outline

- 1 Optics and two-particle dynamics: Why BSE?
- 2 The Bethe-Salpeter equation: Pictorial derivation
- 3 Macroscopic response and the Bethe-Salpeter equation**
- 4 The Bethe-Salpeter equation in practice

# Response to external potential


$$V^{ext} \mapsto n^{ind} \mapsto V^{ind}(\mathbf{r}) = \int d\mathbf{r}' v(\mathbf{r} - \mathbf{r}') n^{ind}(\mathbf{r}') = v n^{ind}$$

Total field acting on particles in the system :  $V^{tot} = V^{ext} + V^{ind}$

## Linear response theory: Definition of the dielectric function

$$n^{ind}(1) = \int d2 \chi(12) V^{ext}(2) \quad \mapsto \quad V^{tot} = (1 + v\chi) V^{ext} \equiv \epsilon^{-1} V^{ext}$$

The density response function  $\chi(12)$  is related to the e-h propagator  $L$



$$\chi(12) = \chi(\mathbf{r}_1, \mathbf{r}_2, t_1 - t_2) = L(1122) = L(\mathbf{r}_1 \mathbf{r}_1 \mathbf{r}_2 \mathbf{r}_2, t_1 - t_2)$$



# Macroscopic response in solids

Optical absorption is determined by  $\text{Im}\epsilon_M(\omega)$ . How we calculate it?

$$V^{ext}(\mathbf{r}, t) = V^{ext}(\mathbf{q})e^{-i(\omega t - \mathbf{q}\mathbf{r})}, \quad q \ll G$$

In a periodic system  $V^{ind}$  contains all components with  $\mathbf{k} = \mathbf{q} + \mathbf{G}$

$$V^{ind}(\mathbf{r}, t) = e^{-i\omega t} \sum_{\mathbf{G}} V_{\mathbf{G}}^{ind}(\mathbf{q}) e^{i(\mathbf{q} + \mathbf{G})\mathbf{r}}$$

Fourier component of the total potential in a solid:

$$V_{\mathbf{G}}^{tot}(\mathbf{q}) = \delta_{\mathbf{G},0} V^{ext}(\mathbf{q}) + V_{\mathbf{G}}^{ind}(\mathbf{q}) = [\delta_{\mathbf{G},0} + v_{\mathbf{G}}(\mathbf{q})\chi_{\mathbf{G},0}(\mathbf{q}, \omega)] V^{ext}(\mathbf{q})$$

## Macroscopic field and macroscopic dielectric function

- Macroscopic (averaged) potential:  $V_M^{tot}(\mathbf{q}) = V_{\mathbf{G}=0}^{tot}(\mathbf{q})$
- Macroscopic dielectric function:  $V^{ext}(\mathbf{q}) = \epsilon_M(\mathbf{q}, \omega) V_M^{tot}(\mathbf{q})$

$$\epsilon_M(\mathbf{q}, \omega) = \frac{1}{1 + v_{\mathbf{G}=0}(\mathbf{q})\chi_{0,0}(\mathbf{q}, \omega)}$$

# Macroscopic dielectric function from BSE (1)

## 1st possibility:

- Calculate  $L(1234)$  by solving the Bethe-Salpeter equation

$$L = L_0 + L_0(v - \gamma)L$$

- Join electron-hole ends and perform a Fourier transform in time

$$L(1122) = L(\mathbf{r}_1\mathbf{r}_1\mathbf{r}_2\mathbf{r}_2, t_1 - t_2) \mapsto L(\mathbf{r}_1\mathbf{r}_1\mathbf{r}_2\mathbf{r}_2, \omega) = \chi(\mathbf{r}_1, \mathbf{r}_2, \omega)$$

- Go to the momentum representation

$$\chi_{\mathbf{G}, \mathbf{G}'}(\mathbf{q}, \omega) = \int d\mathbf{r}_1 d\mathbf{r}_2 e^{i(\mathbf{q} + \mathbf{G})\mathbf{r}_1} L(\mathbf{r}_1\mathbf{r}_1\mathbf{r}_2\mathbf{r}_2, \omega) e^{-i(\mathbf{q} + \mathbf{G}')\mathbf{r}_2}$$

- The “head” of  $\chi_{\mathbf{G}, \mathbf{G}'}$  (element with  $\mathbf{G} = \mathbf{G}' = 0$ ) determines  $\epsilon_M$

# Macroscopic dielectric function from BSE (1)

## 1st possibility:

- Calculate  $L(1234)$  by solving the Bethe-Salpeter equation

$$L = L_0 + L_0(v - \gamma)L$$

- Join electron-hole ends and perform a Fourier transform in time

$$L(1122) = L(\mathbf{r}_1\mathbf{r}_1\mathbf{r}_2\mathbf{r}_2, t_1 - t_2) \mapsto L(\mathbf{r}_1\mathbf{r}_1\mathbf{r}_2\mathbf{r}_2, \omega) = \chi(\mathbf{r}_1, \mathbf{r}_2, \omega)$$

- Go to the momentum representation

$$\chi_{\mathbf{G}, \mathbf{G}'}(\mathbf{q}, \omega) = \int d\mathbf{r}_1 d\mathbf{r}_2 e^{i(\mathbf{q} + \mathbf{G})\mathbf{r}_1} L(\mathbf{r}_1\mathbf{r}_1\mathbf{r}_2\mathbf{r}_2, \omega) e^{-i(\mathbf{q} + \mathbf{G}')\mathbf{r}_2}$$

- The “head” of  $\chi_{\mathbf{G}, \mathbf{G}'}$  (element with  $\mathbf{G} = \mathbf{G}' = 0$ ) determines  $\epsilon_M$

## Macroscopic dielectric function and the absorption rate

$$\epsilon_M(\mathbf{q}, \omega) = \frac{1}{1 + v_{\mathbf{G}=0}(\mathbf{q})\chi_{0,0}(\mathbf{q}, \omega)}; \quad Abs(\omega) = \lim_{\mathbf{q} \rightarrow 0} \epsilon''_M(\mathbf{q}, \omega)$$

# Macroscopic dielectric function from BSE (2)

## 2nd possibility:

Define a “long-range part”  $v_0$  of the interaction potential

$$v_{\mathbf{G}}(\mathbf{q}) = v_{\mathbf{G}=0}(\mathbf{q})\delta_{\mathbf{G},0} + \bar{v}_{\mathbf{G}}(\mathbf{q})$$

$$v(r) = \int_{BZ} d\mathbf{q} \sum_{\mathbf{G}} e^{i(\mathbf{q}+\mathbf{G})\mathbf{r}} v_{\mathbf{G}}(\mathbf{q}) = v_0(\mathbf{r}) + \bar{v}(\mathbf{r})$$

Bethe-Salpeter equation for a “proper” e-h propagator  $\bar{L}$  (1234)  
(replace  $v \mapsto \bar{v}$  in the full BSE)

$$\bar{L} = L_0 + L_0(\bar{v} - \gamma)\bar{L}$$

The full  $L$ -function and the density response function  $\chi_{\mathbf{G},\mathbf{G}'}(\mathbf{q},\omega)$

$$L = \bar{L} + \bar{L}v_0L \quad \Rightarrow \quad \chi = \bar{\chi} + \bar{\chi}v_0\chi$$

# Macroscopic dielectric function from BSE (2)

$$L = \bar{L} + \bar{L}v_0L \quad \Rightarrow \quad \chi(12) = \bar{\chi}(12) + \bar{\chi}(13)v_0(34)\chi(42)$$

In the momentum representation  $v_0 \mapsto v_{\mathbf{G}=0}(\mathbf{q})\delta_{\mathbf{G},0}$

$$\chi_{\mathbf{G},\mathbf{G}'} = \bar{\chi}_{\mathbf{G},\mathbf{G}'} + \bar{\chi}_{\mathbf{G},0}v_{\mathbf{G}=0}\chi_{0,\mathbf{G}'} \quad \Rightarrow \quad \chi_{0,0}(\mathbf{q},\omega) = \frac{\bar{\chi}_{0,0}(\mathbf{q},\omega)}{1 - v_{\mathbf{G}=0}(\mathbf{q})\bar{\chi}_{0,0}(\mathbf{q},\omega)}$$

## Macroscopic dielectric function in terms of proper polarizability

$$\epsilon_M(\mathbf{q},\omega) = \frac{1}{1 + v_{\mathbf{G}=0}(\mathbf{q})\chi_{0,0}(\mathbf{q},\omega)} = 1 - v_{\mathbf{G}=0}(\mathbf{q})\bar{\chi}_{0,0}(\mathbf{q},\omega)$$

$$\bar{\chi}_{0,0}(\mathbf{q},\omega) = \int d\mathbf{r}_1 d\mathbf{r}_2 e^{i\mathbf{q}(\mathbf{r}_1 - \mathbf{r}_2)} \bar{L}(\mathbf{r}_1 \mathbf{r}_1 \mathbf{r}_2 \mathbf{r}_2, \omega)$$

# Macroscopic dielectric function from BSE (2)

## Optical response from the Bethe-Salpeter equation

- Solve the reduced Bethe-Salpeter equation for  $\bar{L}$ (1234)

$$\bar{L} = L_0 + L_0(\bar{v} - \gamma)\bar{L}$$

- Calculate the macroscopic dielectric function from  $\bar{L}$ (1122)

$$\epsilon_M(\mathbf{q}, \omega) = 1 - v_{\mathbf{G}=0}(\mathbf{q}) \int d\mathbf{r}_1 d\mathbf{r}_2 e^{i\mathbf{q}(\mathbf{r}_1 - \mathbf{r}_2)} \bar{L}(\mathbf{r}_1 \mathbf{r}_1 \mathbf{r}_2 \mathbf{r}_2, \omega)$$

- Calculate the absorption rate from the imaginary part of  $\epsilon_M(\mathbf{q}, \omega)$

$$Abs(\omega) = \lim_{\mathbf{q} \rightarrow 0} \epsilon_M''(\mathbf{q}, \omega)$$

By setting  $\bar{v} = 0$  we neglect local field effects – the difference between the macroscopic field  $V_M^{tot}(\mathbf{r})$  and the actual field  $V^{tot}(\mathbf{r})$

# Outline

- 1 Optics and two-particle dynamics: Why BSE?
- 2 The Bethe-Salpeter equation: Pictorial derivation
- 3 Macroscopic response and the Bethe-Salpeter equation
- 4 The Bethe-Salpeter equation in practice**


# The Bethe-Salpeter equation: Approximations

## Reminder


BSE determines 2-particle propagator  $L(1234)$ , provided 1-particle self-energy  $\Sigma(12)$  and e-h scattering amplitude  $\gamma(1234)$  are given.

### Standard approximations:

- Approximating  $\Sigma$  by GW diagram:  $\Sigma(12) = G(12)W(12)$



- Approximating  $\gamma$  by  $W$ :  $\gamma(1234) = W(12)\delta(13)\delta(24)$





# The Bethe-Salpeter equation: Approximations

## Approximate Bethe-Salpeter equation



## Analytic form of the approximate Bethe-Salpeter equation

$$L(1234) = L_0(1234) + \int L_0(1256)[v(57)\delta(56)\delta(78) - W(56)\delta(57)\delta(68)]L(7834)d5d6d7d8$$

$L_0(1234) = G(12)G(43)$  and  $W(12)$  come out of the GW calculations

# The Bethe-Salpeter equation: Approximations

## Reduced BSE for the proper e-h propagator

$$L(1234) = L_0(1234) + \int d5d6d7d8 L_0(1256) \times \\ \times [v(57)\delta(56)\delta(78) - W(56)\delta(57)\delta(68)]L(7834)$$

Further simplifications: Static  $W$

Assumption of the static screening:

$$W(\mathbf{r}_1, \mathbf{r}_2, t_1 - t_2) \Rightarrow W(\mathbf{r}_1, \mathbf{r}_2)\delta(t_1 - t_2)$$

$$\bar{L}(1234) \Rightarrow \bar{L}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, t - t') \Rightarrow \bar{L}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, \omega)$$

# The Bethe-Salpeter equation: Approximations

## Reduced BSE for the proper e-h propagator

$$\begin{aligned} \bar{L}(1234) = & L_0(1234) + \int d5d6d7d8 L_0(1256) \times \\ & \times [\bar{v}(57)\delta(56)\delta(78) - W(56)\delta(57)\delta(68)] \bar{L}(7834) \end{aligned}$$

Further simplifications: Static  $W$

Assumption of the static screening:

$$W(\mathbf{r}_1, \mathbf{r}_2, t_1 - t_2) \Rightarrow W(\mathbf{r}_1, \mathbf{r}_2)\delta(t_1 - t_2)$$

$$\bar{L}(1234) \Rightarrow \bar{L}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, t - t') \Rightarrow \bar{L}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, \omega)$$

# The Bethe-Salpeter equation: Approximations

## Reduced BSE for the proper e-h propagator

$$\bar{L}(1234) = L_0(1234) + \int d5d6d7d8 L_0(1256) \times \\ \times [\bar{v}(57)\delta(56)\delta(78) - W(56)\delta(57)\delta(68)]\bar{L}(7834)$$

### Further simplifications: Static $W$

Assumption of the static screening:

$$W(\mathbf{r}_1, \mathbf{r}_2, t_1 - t_2) \Rightarrow W(\mathbf{r}_1, \mathbf{r}_2)\delta(t_1 - t_2)$$

$$\bar{L}(1234) \Rightarrow \bar{L}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, t - t') \Rightarrow \bar{L}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, \omega)$$

# Optical response in practice

## Calculation of the macroscopic dielectric function

$$\begin{aligned} \bar{L}(\mathbf{r}_1\mathbf{r}_2\mathbf{r}_3\mathbf{r}_4\omega) &= L_0(\mathbf{r}_1\mathbf{r}_2\mathbf{r}_3\mathbf{r}_4\omega) + \int d\mathbf{r}_5d\mathbf{r}_6d\mathbf{r}_7d\mathbf{r}_8 L_0(\mathbf{r}_1\mathbf{r}_2\mathbf{r}_5\mathbf{r}_6\omega) \times \\ &\times [\bar{v}(\mathbf{r}_5\mathbf{r}_7)\delta(\mathbf{r}_5\mathbf{r}_6)\delta(\mathbf{r}_7\mathbf{r}_8) - W(\mathbf{r}_5\mathbf{r}_6)\delta(\mathbf{r}_5\mathbf{r}_7)\delta(\mathbf{r}_6\mathbf{r}_8)] \bar{L}(\mathbf{r}_7\mathbf{r}_8\mathbf{r}_3\mathbf{r}_4\omega) \end{aligned}$$

$$\epsilon_M(\omega) = 1 - \lim_{\mathbf{q} \rightarrow 0} \left[ v_{\mathbf{G}=0}(\mathbf{q}) \int d\mathbf{r}d\mathbf{r}' e^{i\mathbf{q}(\mathbf{r}-\mathbf{r}')} \bar{L}(\mathbf{r}, \mathbf{r}, \mathbf{r}', \mathbf{r}', \omega) \right]$$

$$L_0(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4, \omega) = \sum_{ij} (f_j - f_i) \frac{\phi_i^*(\mathbf{r}_1)\phi_j(\mathbf{r}_2)\phi_i(\mathbf{r}_3)\phi_j^*(\mathbf{r}_4)}{\omega - (E_i - E_j)}$$

# BSE calculations

## A three-step method

### 1 LDA calculation

⇒ Kohn-Sham wavefunctions  $\varphi_i$

### 2 GW calculation

⇒ GW energies  $E_i$  and screened Coulomb interaction  $W$

### 3 BSE calculation


solution of  $\bar{L} = L_0 + L_0(\bar{v} - \gamma)\bar{L}$

⇒ proper e-h propagator  $\bar{L}(\mathbf{r}_1\mathbf{r}_2\mathbf{r}_3\mathbf{r}_4\omega)$

⇒ spectra  $\epsilon_M(\omega)$

# Results: Continuum excitons (Si)


## Bulk silicon



G. Onida, L. Reining, and A. Rubio, RMP **74** (2002).

# Results: Bound excitons (solid Ar)

## Solid argon



F. Sottile, M. Marsili, V. Olevano, and L. Reining, PRB **76** (2007).



# References

## Review articles:



Giovanni Onida, Lucia Reining, and Angel Rubio  
Rev. Mod. Phys. **74**, 601 (2002).



G. Strinati  
Rivista del Nuovo Cimento **11**, (12)1 (1988).



S. Botti, A. Schindlmayr, R. Del Sole, and L. Reining  
Rep. Progr. Phys. **70**, 357 (2007).

## PhD theses:



Francesco Sottile

PhD thesis, Ecole Polytechnique (2003)

[http://etsf.polytechnique.fr/system/files/users/francesco/Tesi\\_dot.pdf](http://etsf.polytechnique.fr/system/files/users/francesco/Tesi_dot.pdf)



Fabien Bruneval

PhD thesis, Ecole Polytechnique (2005)

[http://theory.polytechnique.fr/people/bruneval/bruneval\\_these.pdf](http://theory.polytechnique.fr/people/bruneval/bruneval_these.pdf)