

Exercise 1: Covariant derivative

Prove that the term $\bar{\Psi} \not{D} \Psi$ where the covariant derivative is given by:

$$D_\mu = \partial_\mu - ig\tilde{W}_\mu, \quad \tilde{W}_\mu = T_a W_\mu^a$$

is invariant under gauge transformations:

$$\begin{aligned} \Psi &\mapsto U\Psi, \quad U = \exp\{-iT_a\theta^a(x)\} \\ \tilde{W}_\mu &\mapsto U\tilde{W}_\mu U^\dagger - \frac{i}{g}(\partial_\mu U)U^\dagger \end{aligned}$$

Exercise 2: Non abelian gauge transformations

The Yang-Mills Lagrangian is

$$\mathcal{L}_{\text{YM}} = -\frac{1}{2}\text{Tr}\{\tilde{W}_{\mu\nu}\tilde{W}^{\mu\nu}\}$$

where

$$\tilde{W}_{\mu\nu} \equiv T_a W_{\mu\nu}^a = D_\mu \tilde{W}_\nu - D_\nu \tilde{W}_\mu = \partial_\mu \tilde{W}_\nu - \partial_\nu \tilde{W}_\mu - ig[\tilde{W}_\mu, \tilde{W}_\nu], \quad \tilde{W}_\mu \equiv T_a W_\mu^a$$

and T_a are the N generators of a Lie group with algebra $[T_a, T_b] = if_{abc}T_c$.

i) Check that under a gauge transformation of the fields:

$$\tilde{W}_\mu \mapsto U\tilde{W}_\mu U^\dagger - \frac{i}{g}(\partial_\mu U)U^\dagger, \quad U = \exp\{-iT_a\omega^a\}$$

the $\tilde{W}_{\mu\nu}$ transforms as

$$\tilde{W}_{\mu\nu} \mapsto U\tilde{W}_{\mu\nu}U^\dagger$$

and therefore \mathcal{L}_{YM} is gauge invariant.

ii) Check that one may write

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4}W_{\mu\nu}^a W^{a,\mu\nu}$$

that contains kinetic terms and cubic and quartic interactions among the gauge fields.

iii) Check that


$$W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + gf_{abc}W_\mu^b W_\nu^c$$

iv) Check that under infinitesimal gauge transformations:

$$W_\mu^a \mapsto W_\mu^a - f_{abc}W_\mu^b \omega^c - \frac{1}{g}\partial_\mu \omega^a$$


Exercise 3: Feynman rules of general non-Abelian gauge theories

Obtain the Feynman rules for cubic and quartic self-interactions among gauge fields in a general non-Abelian gauge theory, as well as those for the interactions of Faddeev-Popov ghosts with gauge fields:



Exercise 4: Faddeev-Popov ghosts and gauge invariance

Consider the 1-loop self-energy diagrams for non-Abelian gauge theories in the figure. Calculate the diagrams in the 't Hooft-Feynman gauge and show that the sum does not have the tensor structure $g_{\mu\nu}k^2 - k_\mu k_\nu$ required by the gauge invariance of the theory unless diagram (c) involving ghost fields is included.



Hint: Take Feynman rules from previous exercise and use dimensional regularization. It is convenient to use the Passarino-Veltman tensor decomposition of loop integrals:

$$\frac{i}{16\pi^2} \{B_0, B_\mu, B_{\mu\nu}\} = \mu^\epsilon \int \frac{d^D q}{(2\pi)^D} \frac{\{1, q_\mu, q_\mu q_\nu\}}{q^2(q+k)^2}$$

where $B_0 = \Delta_\epsilon + \text{finite}$

$$B_\mu = k_\mu B_1, \quad B_1 = -\frac{\Delta_\epsilon}{2} + \text{finite}$$

$$B_{\mu\nu} = g_{\mu\nu} B_{00} + k_\mu k_\nu B_{11}, \quad B_{00} = -\frac{k^2}{12} \Delta_\epsilon + \text{finite}, \quad B_{11} = \frac{\Delta_\epsilon}{3} + \text{finite}$$

with $\Delta_\epsilon = 2/\epsilon - \gamma + \ln 4\pi$ and $D = 4 - \epsilon$. You may check that the ultraviolet divergent part has the expected structure or find the final result in terms of scalar integrals, that for massless fields read:

$$B_1 = -\frac{1}{2} B_0, \quad B_{00} = -\frac{k^2}{4(D-1)} B_0, \quad B_{11} = \frac{D}{4(D-1)} B_0.$$

Do not forget a symmetry factor (1/2) in front of (a) and (b), and a factor (-1) in (c).

Exercise 5: Propagator of a massive vector boson field

Consider the Proca Lagrangian of a massive vector boson field

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}M^2A_\mu A^\mu, \quad \text{with} \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu.$$

Show that the propagator of A_μ is

$$\tilde{D}_{\mu\nu}(k) = \frac{i}{k^2 - M^2 + i0} \left[-g_{\mu\nu} + \frac{k_\mu k_\nu}{M^2} \right]$$

Exercise 6: Propagator of a massive gauge field

Consider the U(1) gauge invariant Lagrangian \mathcal{L} with gauge fixing \mathcal{L}_{GF} :

$$\begin{aligned} \mathcal{L} &= -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu\phi)^\dagger(D^\mu\phi) - \mu^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2 \\ \mathcal{L}_{\text{GF}} &= -\frac{1}{2\xi}(\partial_\mu A^\mu - \xi M_A\chi)^2, \quad \text{with} \quad D_\mu = \partial_\mu + ieA_\mu, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \end{aligned}$$

where $M_A = ev$ after spontaneous symmetry breaking ($\mu^2 < 0$, $\lambda > 0$) when the complex scalar field ϕ acquires a VEV and is parameterized by

$$\phi(x) = \frac{1}{\sqrt{2}}[v + \varphi(x) + i\chi(x)], \quad \mu^2 = -\lambda v^2.$$

Show that the propagators of φ , χ and the gauge field A_μ are respectively

$$\begin{aligned} \tilde{D}^\varphi(k) &= \frac{i}{k^2 - M_\varphi^2 + i0} \quad \text{with} \quad M_\varphi^2 = -2\mu^2 = 2\lambda v^2 \\ \tilde{D}^\chi(k) &= \frac{i}{k^2 - \xi M_A^2 + i0}, \quad \tilde{D}_{\mu\nu}(k) = \frac{i}{k^2 - M_A^2 + i0} \left[-g_{\mu\nu} + (1 - \xi)\frac{k_\mu k_\nu}{k^2 - \xi M_A^2} \right] \end{aligned}$$

Exercise 7: The conjugate Higgs doublet

Show that $\Phi^c \equiv i\sigma_2\Phi^* = \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix}$ transforms under SU(2) like $\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$, with $\phi^- = (\phi^+)^*$. What are the weak isospins, hypercharges and electric charges of ϕ^0 , ϕ^{0*} , ϕ^+ , ϕ^- ?
Hint: Use the property of Pauli matrices: $\sigma_i^* = -\sigma_2\sigma_i\sigma_2$.

Exercise 8: Lagrangian and Feynman rules of the Standard Model

Try to reproduce the Lagrangian and the corresponding Feynman rules of as many Standard Model interactions as you can. Of particular interest/difficulty are [VVV] and [VVVV].

Exercise 9: Z pole observables at tree level

Show that

$$(a) \quad \Gamma(f\bar{f}) \equiv \Gamma(Z \rightarrow f\bar{f}) = N_c^f \frac{\alpha M_Z}{3} (v_f^2 + a_f^2), \quad N_c^f = 1 \text{ (3) for } f = \text{lepton (quark)}$$

$$(b) \quad \sigma_{\text{had}} = 12\pi \frac{\Gamma(e^+e^-)\Gamma(\text{had})}{M_Z^2 \Gamma_Z^2}$$

$$(c) \quad A_{FB} = \frac{3}{4} A_f, \quad \text{with } A_f = \frac{2v_f a_f}{v_f^2 + a_f^2}$$

Exercise 10: Higgs partial decay widths at tree level

Show that

$$(a) \quad \Gamma(H \rightarrow f\bar{f}) = N_c^f \frac{G_F M_H}{4\pi\sqrt{2}} m_f^2 \left(1 - \frac{4m_f^2}{M_H^2}\right)^{3/2}, \quad N_c^f = 1 \text{ (3) for } f = \text{lepton (quark)}$$

$$(b) \quad \Gamma(H \rightarrow W^+W^-) = \frac{G_F M_H^3}{8\pi\sqrt{2}} \sqrt{1 - \frac{4M_W^2}{M_H^2}} \left(1 - \frac{4M_W^2}{M_H^2} + \frac{12M_W^4}{M_H^4}\right)$$

$$\Gamma(H \rightarrow ZZ) = \frac{G_F M_H^3}{16\pi\sqrt{2}} \sqrt{1 - \frac{4M_Z^2}{M_H^2}} \left(1 - \frac{4M_Z^2}{M_H^2} + \frac{12M_Z^4}{M_H^4}\right)$$