

MINIMAL KITAEV-TRANSMON QUBIT BASED ON DOUBLE QUANTUM DOTS

D. Michel Pino¹
 T: + 34 675 84 50 82, dmichel.pino@csic.es
 Rubén Seoane Souto¹
 Ramón Aguado¹

¹Instituto de Ciencia de Materiales de Madrid (ICMM), Consejo Superior de Investigaciones Científicas (CSIC), Sor Juana Inés de la Cruz 3, 28049 Madrid, Spain

Minimal Kitaev chains composed of two semiconducting quantum dots coupled via a superconductor have emerged as a promising platform to realize and study Majorana bound states (MBSs), which appear for fine-tuned configurations. We propose a hybrid qubit based on a Josephson junction between two such double quantum dots (DQDs) embedded in a superconducting qubit geometry. The qubit makes use of the 4π -Josephson effect in the Kitaev junction to create a subspace based on the even/odd fermionic parities of the two DQD arrays hosting MBSs. Deep in the transmon regime, we demonstrate that by performing circuit QED spectroscopy on such a hybrid Kitaev-transmon “Kitmon” qubit, one could observe distinct MBS features in perfect agreement with precise analytical predictions in terms of DQD parameters only. This agreement allows us to extract the Majorana polarization in the junction from the microwave response.




Figure 1 – Schematic illustration of the Kitaev-transmon device.




Figure 2 – Majorana polarization and Majorana coupling.

References:

- [1] T. Dvir et al., Nature 614, 7948 (2023) 445.
- [2] M. Leijnse and K. Flensberg, Phys. Rev. B 86, 13 (2012) 134528.
- [3] D. Michel Pino, Rubén Seoane Souto and Ramón Aguado, Phys. Rev. B 109, 7 (2024) 075101.




Figure 3 – Kitaev-transmon qubit spectroscopy.